



---

**Clinical Study Protocol**

Drug Substance     Dapagliflozin  
Study Code         D1692C00012  
Edition Number     1  
Date

---

---

**A long term open label study to evaluate the safety and efficacy of dapagliflozin as monotherapy or combination therapies with anti-diabetic drugs in Japanese subjects with type 2 diabetes who have inadequate glycemic control**

---

**Sponsor:**

AstraZeneca K.K.  
Bristol-Myers K.K.

**The following Amendment(s) and Administrative Changes have been made to this protocol since the date of preparation:**

<b>Amendment No.</b>	<b>Date of Amendment</b>	<b>Local Amendment No:</b>	<b>Date of Local Amendment</b>
_____	_____	_____	_____
_____	_____	_____	_____
_____	_____	_____	_____
<b>Administrative Change No.</b>	<b>Date of Administrative Change</b>	<b>Local Administrative Change No.</b>	<b>Date of Local Administrative Change</b>
_____	_____	_____	_____
_____	_____	_____	_____
_____	_____	_____	_____

This submission /document contains trade secrets and confidential commercial information, disclosure of which is prohibited without providing advance notice to Bristol-Myers Squibb and AstraZeneca and opportunity to object.

## PROTOCOL SYNOPSIS

---

### **A long term open label study to evaluate the safety and efficacy of dapagliflozin as monotherapy or combination therapies with anti-diabetic drugs in Japanese subjects with type 2 diabetes who have inadequate glycemic control**

---

#### **Principal Investigator**

The names of the investigators are described in Supplement A, “Investigators and Study Administrative Structure”.

#### **Study centre(s) and number of subjects planned**

**Study Centres:** approximately 100 centres are planned for participation

**Number of Subjects:** 700 treated Japanese subjects

---

<b>Study period</b>	<b>Phase of development</b>	
Estimated date of first subject enrolled	February 2011	Phase III
Estimated date of last subject completed	October 2012	

---

#### **Objectives**

##### **Primary Objective:**

To evaluate safety and tolerability of long term treatment up to 52 weeks with the dosing regimen of dapagliflozin, where it starts with 5 mg and titrates up to 10 mg depending on subjects' condition of glycemic control, for total subjects and for each subgroup of monotherapy or combination therapies with any anti-diabetic drugs including sulphonylureas (SU), rapid-acting insulin secretagogue (glinides), metformin,  $\alpha$ -glucosidase inhibitors (AGI), thiazolidinedions (TZD), dipeptidyl peptidase-4 inhibitors (DPP-4) and glucagon-like peptide-1 agonists (GLP-1).

##### **Secondary Objectives:**

To evaluate the efficacy of the dosing regimen of dapagliflozin over period of time up to 52 weeks for each subgroup of monotherapy or combination therapy with SU, glinides, metformin, AGI, TZD, DPP-4 or GLP-1 with regard to glycemic, weight and blood pressure (BP) parameters.

## **Study design**

This is a long term, single arm, open label study to evaluate the safety and efficacy of dapagliflozin as monotherapy or combination therapies with anti-diabetic drugs including SU, glinides, metformin, AGI, TZD, DPP-4 and GLP-1 in Japanese subjects of type 2 diabetes mellitus (T2DM) who have inadequate glycaemic control.

## **Target subject population**

Total 700 Japanese subjects with T2DM who have inadequate glycaemic control on diet and exercise or on concomitant anti-diabetic treatment

### Subgroups:

- Monotherapy: 240 subjects. The treatment-naïve subjects and the subjects who received ongoing medical treatment for diabetes within 6 weeks of enrolment, except for TZD (single oral anti-diabetic agent or two agents with less than half of the approved maximal dose for each agent) need a wash-out period.
- Combination therapies with SUs: 120 subjects. 8 weeks or more treatment before start of dapagliflozin treatment is needed.
- Combination therapies with DPP-4, AGI, metformin, or TZD: 60 subjects for each. 8 weeks or more treatment for AGI, metformin, DPP-4, or 12 weeks or more for TZD before start of dapagliflozin treatment is needed.
- Combination therapy with glinides or GLP-1: 50 subjects for each. 8 weeks or more treatment before start of dapagliflozin treatment is needed.

Either gender needs to be 40% or higher of total number of treated subjects for each sub-group. Only in the monotherapy group, the proportion of subjects with Haemoglobin A1c (HbA1c)  $\geq 6.5\%$  but  $\leq 7\%$  at Visit 5 needs to be at most approximately 25%.

## **Investigational product, dosage and mode of administration**

Investigational product should be taken once daily in the morning.

- Dapagliflozin 5 mg dose: Dapagliflozin 5 mg 1 tablet
- Dapagliflozin 10 mg dose: Dapagliflozin 10 mg 1 tablet

## **Comparator, dosage and mode of administration**

Not applicable

## **Duration of treatment**

In the monotherapy subgroup, this study consists of a 2-week screening period, a 4-week lead-in period, a 52-week treatment period and a 3-week follow-up period. The subjects who

received ongoing medical treatment for diabetes within 6 weeks of enrolment, except for TZD (a single oral anti-diabetic agent or two agents with less than half of the approved maximal dose for each agent) need a 6-week wash-out period before a 4-week lead-in period.

In the combination therapy subgroup, this study consists of a 2-week screening period, a 4-week lead-in period, a 52-week treatment period and a 3-week follow-up period.

The total planned study duration starting with enrolment and including the follow-up period will be 61 weeks.

Dapagliflozin starts with 5 mg at Visit 6 (Week 0) and is up-titrated to 10 mg at any visit after Week 12 (Visit 10) under the conditions of HbA1c >7.5% and no safety concerns.

**Outcome variable(s):**

**Primary outcome variables (Safety):**

- Adverse Events (AEs) and Serious adverse events (SAEs)
- Hypoglycemic events
- Safety laboratory values
- Electrocardiogram (ECG)
- Vital Signs (pulse and BP)
- Physical examination

**Statistical methods**

Descriptive summaries of demographics and baseline characteristics will be presented by subgroup and for all subjects combined using the safety analysis set. Descriptive statistics at each time point, including Week 52 (LOCF) will be presented by subgroup as well as for total population for all efficacy variables. Safety variables will be summarized descriptively.

	<b>PAGE</b>
TITLE PAGE .....	1
PROTOCOL SYNOPSIS.....	2
TABLE OF CONTENTS.....	5
LIST OF ABBREVIATIONS AND DEFINITION OF TERMS .....	11
1. INTRODUCTION .....	13
1.1 Background .....	13
1.2 Research hypothesis.....	14
1.3 Rationale for conducting this study .....	14
1.4 Benefit/risk and ethical assessment.....	15
2. STUDY OBJECTIVES.....	17
2.1 Primary objective .....	17
2.2 Secondary objectives .....	17
2.3 Safety objective.....	17
2.4 Exploratory objectives – Not applicable.....	18
3. STUDY PLAN AND PROCEDURES .....	18
3.1 Overall study design and flow chart .....	18
3.2 Monotherapy subgroup .....	19
3.2.1 Screening Period (2 weeks) : Visit 1-2 .....	19
3.2.2 Wash-out period (6 weeks) : Visit 2-4.....	19
3.2.3 Lead-in Period (4 weeks) :Visit 4-6.....	20
3.2.4 Treatment Period (52 weeks):Visit 6-17.....	20
3.2.4.1 Rescue Treatment.....	21
3.2.5 Follow-up period (3 weeks) : Visit 17-18.....	21
3.3 Combination therapy subgroup.....	22
3.3.1 Screening Period (2 weeks) : Visit 1 and 4.....	22
3.3.2 Lead-in Period (4 weeks) : Visit 4-6.....	22
3.3.3 Treatment Period (52 weeks) : Visit 6-17 .....	23
3.3.3.1 Criteria for reduction of basal SU.....	23
3.3.3.2 Rescue Treatment.....	23
3.3.4 Follow-up period (3 weeks) : Visit 17-18.....	24
3.4 Rationale for study design, doses and control groups.....	31
3.4.1 Study design and regulatory requirement .....	31

3.4.2	Study doses and control groups .....	31
3.4.3	Choice of outcome variables.....	32
3.4.4	Choice of study population .....	32
4.	SUBJECT SELECTION CRITERIA .....	33
4.1	Inclusion criteria .....	33
4.2	Exclusion criteria .....	35
5.	STUDY CONDUCT .....	39
5.1	Restrictions during the study .....	39
5.2	Subject enrolment .....	39
5.3	Procedures for handling subjects incorrectly enrolled or treated.....	40
5.4	Blinding and procedures for unblinding the study – Not applicable .....	40
5.5	Treatments.....	40
5.5.1	Identity of investigational product(s).....	40
5.5.2	Doses and treatment regimens .....	41
5.5.3	Additional study drug – Not applicable .....	41
5.5.4	Labelling .....	41
5.5.5	Storage .....	41
5.6	Concomitant and post-study treatment(s) .....	41
5.7	Treatment compliance.....	42
5.7.1	Accountability.....	43
5.8	Discontinuation of investigational product.....	43
5.8.1	Criteria for discontinuation from the study.....	43
5.8.2	Procedures for discontinuation of a subject from investigational product .....	45
5.9	Withdrawal from study .....	46
6.	COLLECTION OF STUDY VARIABLES.....	46
6.1	Recording of data.....	46
6.2	Data collection and enrolment .....	46
6.2.1	Follow-up procedures .....	46
6.3	Efficacy .....	47
6.3.1	Efficacy laboratory variables .....	47
6.3.2	HbA1c .....	47
6.3.3	FPG .....	47
6.3.4	Weight and height .....	47
6.3.5	BMI .....	48
6.3.6	Blood pressure .....	48
6.3.7	Waist circumference .....	48
6.4	Safety .....	48
6.4.1	Definition of adverse events .....	48

6.4.2	Definitions of serious adverse event .....	48
6.4.3	Recording of adverse events .....	49
6.4.4	Reporting of serious adverse events.....	52
6.4.5	Laboratory safety assessment .....	53
6.4.6	Physical examination .....	56
6.4.7	ECG.....	56
6.4.8	Vital signs .....	56
6.4.8.1	Pulse and blood pressure.....	56
6.4.8.2	Orthostatic blood pressure .....	57
6.4.9	Other safety assessments.....	57
6.4.9.1	Fasting plasma glucose concentrations and hypoglycemic events .....	57
6.4.9.2	Urinary and Genital Infections.....	59
6.4.10	Congestive Heart Failure .....	61
6.4.11	Change in kidney function .....	61
6.4.12	Hyponatremia.....	61
6.4.13	CK abnormalities .....	61
6.4.14	Liver function test abnormalities .....	61
6.4.15	Independent Adjudication Committee .....	61
6.4.16	Hepatic Adjudication Committee .....	62
6.5	Patient reported outcomes (PRO) – Not applicable.....	63
6.6	Pharmacokinetics – Not applicable.....	63
6.7	Pharmacodynamics – Not applicable.....	63
6.8	Pharmacogenetics – Not applicable .....	63
6.9	Health economics – Not applicable .....	63
7.	BIOLOGICAL SAMPLING PROCEDURES.....	63
7.1	Volume of blood .....	63
7.2	Handling, storage and destruction of biological samples .....	63
7.3	Labelling and shipment of biohazard samples.....	63
7.4	Chain of custody of biological samples .....	63
7.5	Withdrawal of informed consent for donated biological samples .....	64
8.	ETHICAL AND REGULATORY REQUIREMENTS.....	64
8.1	Ethical conduct of the study.....	64
8.2	Subject data protection.....	65
8.3	Ethics and regulatory review.....	65
8.4	Informed consent .....	66
8.5	Changes to the protocol and informed consent form .....	66
8.6	Audits and inspections .....	67

9.	STUDY MANAGEMENT BY ASTRAZENECA .....	67
9.1	Pre-study activities .....	67
9.2	Training of study site personnel .....	67
9.3	Monitoring of the study .....	67
9.3.1	Source data .....	68
9.3.2	Direct access to source data in Japan .....	68
9.4	Study agreements .....	69
9.4.1	Archiving of study documents .....	69
9.4.2	Deviation from the clinical study protocol in Japan .....	70
9.5	Study timetable and end of study .....	70
10.	DATA MANAGEMENT BY ASTRAZENECA DATA MANAGEMENT CENTRE .....	71
11.	EVALUATION AND CALCULATION OF VARIABLES BY ASTRAZENECA .....	72
11.1	Calculation or derivation of efficacy variable(s) .....	72
11.1.1	Change and percent change from baseline .....	72
11.1.2	Last observation carried forward (LOCF) .....	72
11.2	Calculation or derivation of safety variable(s) .....	72
11.2.1	Calculation of eGFR .....	72
11.2.2	Other significant adverse events (OAE) .....	72
11.3	Calculation or derivation of patient reported outcome variables – Not applicable .....	73
11.4	Calculation or derivation of pharmacokinetic variables – Not applicable .....	73
11.5	Calculation or derivation of pharmacodynamic variable(s) – Not applicable .....	73
11.6	Calculation or derivation of pharmacogenetic variables – Not applicable .....	73
11.7	Calculation or derivation of health economic variables – Not applicable .....	73
12.	STATISTICAL METHODS AND SAMPLE SIZE DETERMINATION BY ASTRAZENECA .....	73
12.1	Description of analysis sets .....	73
12.1.1	Full analysis set .....	73
12.1.2	Safety analysis set .....	73
12.2	Methods of statistical analyses .....	74
12.2.1	Demographics and baseline characteristics .....	74
12.2.2	Efficacy analysis .....	74
12.2.3	Safety analysis .....	74
12.3	Determination of sample size .....	74
12.4	Data monitoring committee – Not applicable .....	75



13.	IMPORTANT MEDICAL PROCEDURES TO BE FOLLOWED BY THE INVESTIGATOR .....	75
13.1	Medical emergencies and AstraZeneca contacts .....	75
13.2	Overdose .....	75
13.3	Pregnancy.....	75
14.	LIST OF REFERENCES .....	76

## LIST OF TABLES

Table 1	Summary of subgroups.....	18
Table 2	Criteria for initiation of rescue therapy during the treatment period.....	21
Table 3	Criteria for initiation of rescue therapy during the treatment period.....	24
Table 4	Study plan for subgroup of monotherapy .....	25
Table 5	Study plan for subgroups of combination therapy.....	28
Table 6	Identity of investigational product(s) .....	40
Table 7	Drug dispensing scheme.....	41
Table 8	Efficacy laboratory variables.....	47
Table 9	Safety laboratory variables .....	54
Table 10	Volume of blood to be drawn from each subject (mL) .....	63

## LIST OF FIGURES

Figure 1	Study flow chart (monotherapy).....	19
Figure 2	Study flow chart (combination therapies) .....	22

## LIST OF APPENDICES

Appendix A	Signatures
Appendix B	Additional Safety Information
Appendix C	IATA 6.2 Guidance document
Appendix D	New York Heart Association (NYHA) Classification

Appendix E      Algorithm on Management of Hyponatraemia

Appendix F      Case Identification and Management of Decreased Renal Function

Appendix G      Algorithm on Management of Sustained Elevated Liver Safety Abnormalities

## **LIST OF SUPPLEMENT**

Supplement A      Investigators and Study Administrative Structure

## LIST OF ABBREVIATIONS AND DEFINITION OF TERMS

The following abbreviations and special terms are used in this study Clinical Study Protocol.

<b>Abbreviation or special term</b>	<b>Explanation</b>
AE	Adverse Event (See definition in Section 6.4.1)
AGI	$\alpha$ -Glucosidase Inhibitors
ALT	Alanine aminotransferase
AST	Aspartate aminotransferase
BMI	Body Mass Index
CEC	Clinical Event Committee
CK	Creatinine Kinase
CSA	Clinical Study Agreement
DPP-4	Dipeptidyl Peptidase-4 inhibitors
EC	Ethics Committee, synonymous to Institutional Review Board (IRB) and Independent Ethics Committee (IEC)
ECG	Electrocardiogram
eCRF	electronic Case Report Form
eGFR	estimated Glomerular Filtration Rate
FFA	Free Fatty Acid
FPG	Fasting Plasma Glucose
GCP	Good Clinical Practice
GLP-1	Glucagon-Like Peptide-1 agonists
GMP	Good Manufacturing Practice
HbA1c	Haemoglobin A1c
HCG	Human Chorionic Gonadotropin
HDL-C	High-Density Lipoprotein Cholesterol
IATA	International Air Transport Association
ICH	International Conference on Harmonisation
IRB	Institutional Review Board
IWRS	Interactive Web Response System
LDL-C	Low-Density Lipoprotein Cholesterol
LOCF	Last Observation Carried Forward

---

<b>Abbreviation or special term</b>	<b>Explanation</b>
MedDRA	Medical Dictionary for Regulatory Activities
MHLW	Ministry of Health, Labour and Welfare
MI	Myocardial Infarction
NGSP	National Glycohemoglobin Standardization Program
OAE	Other Significant Adverse Event (See definition in Section <a href="#">11.2.2</a> )
PT	Prothrombin Time
PTH	Parathormone
PTT	Partial Thromboplastin Time
SAE	Serious Adverse Event (See definition in Section <a href="#">6.4.2</a> ).
SCr	Serum Creatinine
SGLT	Sodium-glucose Transporters
SU	Sulphonylureas
TB	Total Bilirubin
TC	Total Cholesterol
TG	Triglyceride
TSH	Thyroid Stimulating Hormone
TZD	Thiazolidinedions
T2DM	Type 2 Diabetes
UACR	Urine Albumin: Creatinine Ratio
ULN	Upper Limit of Normal
UTI	Urinary Tract Infections
WBDC	Web Based Data Capture
WOCBP	Women of Childbearing Potential

---

## 1. INTRODUCTION

### 1.1 Background

Type 2 diabetes (T2DM) is characterised by beta-cell dysfunction and peripheral insulin resistance leading to hyperglycemia ([Matthaei et al 2000](#), [Meier and Butler 2005](#)). Chronic hyperglycemia is associated with the development of both macrovascular (myocardial infarction [MI], stroke, peripheral arterial disease), and microvascular (nephropathy, retinopathy, neuropathy) complications ([UKPDS group 1998](#)). Current treatment regimens aiming to reduce glucose levels in patients with T2DM have focused on the stimulation of insulin secretion (eg, sulphonylureas [SU], glinides, glucagon-like peptide-1 agonists [GLP-1] analogs, dipeptidyl peptidase-4 inhibitors [DPP-4] inhibitors), improvement in insulin sensitivity (eg, metformin, thiazolidinediones [TZD]), inhibition of intestinal glucose absorption (eg, acarbose), or the replacement of insulin. However, the limited efficacy of currently available antihyperglycemic agents, as well as associated side effects (eg, hypoglycemia, oedema, weight gain, etc.) clearly underline the need for novel antihyperglycemic treatment strategies ([ACCORD 2008](#), [Koro et al 2004](#)). In addition, the majority of patients with T2DM require more than one antihyperglycemic agent to achieve glycemic targets ([Nathan 2006](#)).

Intestinal absorption and renal reabsorption of glucose are mediated through sodium-glucose transporters (SGLT) ([Silverman 1991](#)). Two sodium glucose transporters, SGLT1 and SGLT2, have been identified as the major transporters of glucose in humans. SGLT1 is expressed in the gastrointestinal tract, heart, skeletal muscle, liver, lung, and kidney, while SGLT2 is expressed almost exclusively in the kidney ([Silverman 1991](#), [Wright 2001](#)). SGLT2 expression is localized in the S1 segment of the proximal tubule, where according to animal data, >90% of renal glucose reabsorption occurs ([Wright 2001](#)). Thus, SGLT2 appears to be the major transporter responsible for the reabsorption of glucose from the glomerular filtrate.

Human SGLT2 mutations are associated with a condition known as familial renal glucosuria. These individuals have varying degrees of glucosuria; those who have loss of function in both alleles can excrete 100 g of glucose or greater per day. The majority of patients are asymptomatic, and their condition is diagnosed incidentally. Typically they do not have hypoglycemic episodes, electrolyte imbalance or increased risk of urinary tract infections ([Santer et al 2003](#)). Even the most severe form of the condition appears to be associated with a favourable prognosis ([Scholl-Burgi et al 2004](#)) although very few patients have been described in the literature. This human model of SGLT2 inhibition supports the potential safety of this mechanism as a treatment approach for type 2 diabetes by demonstrating that mild to moderate glucosuria in itself is not associated with significant adverse health consequences.

Dapagliflozin is a rationally designed, potent, highly selective and orally active inhibitor of the human renal SGLT2. Dapagliflozin lowers plasma glucose by inhibiting the renal reabsorption of glucose and thereby by promoting its urinary excretion. This compound is being developed as an oral agent for the treatment of T2DM, and represents a novel therapeutic approach for the treatment of this disorder. Proof of concept for dapagliflozin in subjects with T2DM has been established in a global Phase IIb study over a dose range from 2.5 to 50 mg over 12 weeks,

administered orally once daily as monotherapy. In this study, dapagliflozin treatment led to significant and clinically relevant reductions in fasting plasma glucose (FPG), postprandial glucose, and haemoglobin (HbA1c) levels throughout the entire dose range, and was associated with weight loss (Ferrannini, 2010). Similar findings were observed in a global Phase III study with administration of once daily dapagliflozin 2.5 mg, 5 mg and 10 mg as an add-on to metformin in T2DM subjects inadequately controlled on metformin alone (Bailey 2010).

In a Phase IIb study in Japan, similar results were also seen in Japanese subjects with T2DM who had inadequate glycaemic control. A statistically significant mean reduction from baseline in HbA1c was seen in all dapagliflozin treatment groups (1, 2.5, 5 and 10 mg) compared to the placebo group, and a clinically relevant effect on HbA1c (>0.5% placebo corrected reduction) over 12 weeks was seen in the dapagliflozin 5 mg and 10 mg groups. In addition, treatment with dapagliflozin as monotherapy for 12 weeks was safe and well tolerated for Japanese subjects with T2DM and the overall safety profile is similar to other reported dapagliflozin clinical studies with no unexpected findings.

For additional details on the background of dapagliflozin, please see the Investigator's Brochure.

## **1.2 Research hypothesis**

Long term treatment, up to 52 weeks, dapagliflozin is safe and well tolerated as monotherapy or combination therapies with anti-diabetic drugs in Japanese subjects with T2DM who have inadequate glycaemic control.

## **1.3 Rationale for conducting this study**

This study will be performed as part of the clinical development programme for dapagliflozin for the treatment of T2DM.

Dapagliflozin has a new mechanism of action, which results in induction of glucose excretion into urine independent of insulin action, is expected to show an anti-hyperglycaemic effect and lead to caloric loss with associated loss in body weight.

This study intends to investigate the safety and tolerability of dapagliflozin in long-term monotherapy or combination use with one of the hypoglycaemic drugs already approved and marketed for T2DM in the treatment of Japanese subjects who have inadequate glycaemic control.

This study has standard design features for long-term monotherapy and combination therapy studies (eg, multi-center, open label) and incorporates the relevant features of the [Japanese Guideline on Clinical Evaluation methods on Oral Hypoglycaemic Agents](#) which was issued 9<sup>th</sup> July 2010 with regard to duration of treatment, choice of study population, and choice of outcome variables.

## **1.4 Benefit/risk and ethical assessment**

### **Risk category**

Considering dapagliflozin's mechanism of action, the previous clinical experience with dapagliflozin, the study's design features (including the inclusion, exclusion, and discontinuation criteria), and the planned safety procedures, participation in this study presents a minimal and thus acceptable risk to the individual subjects that will be included.

### **Potential risks**

The potential risks associated with dapagliflozin that have been identified based upon the mechanism of action, the preclinical results, and the clinical experience to date, as well as precautions included in the global Phase III programme to monitor and/or minimize these risks, are included in the Investigator Brochure.

Inhibition of SGLT2 results in increased urinary glucose excretions, which is commonly believed to increase the risk of urinary tract infections (UTIs). In some of the global Phase III studies, events suggestive of UTI were reported in a slightly higher proportion of dapagliflozin-treated subjects than the placebo group. Increased urinary glucose excretion may also lead to an increased risk of developing genital infections. In global Phase III studies, the proportions of subjects treated with dapagliflozin who reported adverse events (AEs) that matched a predefined list of Medical dictionary for regulatory activities (MedDRA) preferred terms that were indicative of genital infection were higher than those seen for placebo.

Based on the mechanism of action of dapagliflozin and results of animal and clinical studies, there may be a potential risk for this compound to cause hypovolaemia or electrolyte imbalance. As a precaution, subjects at risk for hypovolaemia or electrolyte disturbance should not receive dapagliflozin until more clinical information is available from human studies. In subjects already receiving dapagliflozin who develop conditions that may cause hypovolaemia or electrolyte imbalance, decisions to interrupt or discontinue dapagliflozin therapy and management of subjects should be based on clinical judgment.

Bone health was evaluated in the dapagliflozin clinical development program due to the possible effects of dapagliflozin on body weight, renal tubular handling of calcium and phosphorus, metabolism of vitamin D. Fractures were reported infrequently and the frequency was similar between the dapagliflozin groups and control in the clinical trials with dapagliflozin. Mean change from baseline in the markers of bone resorption was found to be slightly higher in dapagliflozin-treated subjects compared with placebo-treated subjects. However, change in bone formation markers was inconsistent. Therefore, a definitive conclusion on the net effect of treatment of humans with dapagliflozin on bone turnover (resorption/formation) cannot be made at this time.

Higher proportions of subjects with marked laboratory abnormalities of hyperphosphatemia were reported in dapagliflozin vs. placebo but the clinical meaning of this is unclear.

Hepatic laboratory markers were assessed in all the clinical studies with dapagliflozin. In the pooled analyses, the proportion of subjects with elevated liver function tests was similar in the dapagliflozin and comparator groups and no clinically meaningful or consistent mean changes from baseline in liver function tests were observed in the dapagliflozin and placebo groups across the Phase IIb and III clinical studies. One subject had an serious adverse events (SAE) reported as drug-induced acute hepatitis and was later also diagnosed with probable autoimmune hepatitis.

In addition, all subjects except for monotherapy in this study will receive a long-term combination use of dapagliflozin with broadly categorized classes of the already approved and marketed drugs as background medication. These medications are widely used for anti-diabetic treatments and will be prescribed according to the approved label, however, the safety of the combination therapy has not been established in Japanese subjects.

No study procedure will put subjects at a risk beyond those ordinarily encountered during the performance of routine medical examinations or routine tests.

### **Protection against risks**

This study has been designed with appropriate measures in place so as to monitor and minimize any of the potential health risks to participating subjects. In order to ensure the safety of all subjects participating in this study, AstraZeneca and Bristol-Myers Squibb will conduct a real-time review of all safety information from all ongoing clinical dapagliflozin studies as they become available. Safety signal detection will include the integration of all available sources of safety information, including clinical study data, AE reports, pre-clinical data, epidemiological studies and literature reports, to identify and characterize unrecognized safety risks or changes in those which are currently expected Adverse Drug Reactions. Any information that may affect the benefit-risk profile of dapagliflozin will be immediately communicated to relevant Health Authorities and appropriate actions will be taken regarding the clinical program as needed. Thus real-time, active safety surveillance will be conducted during the entire duration of this study. In addition, all dapagliflozin studies are subject to a carefully designed subject risk management plan that includes the temporary and if necessary permanent discontinuation of investigational product in individual subjects in whom a potential health risk or a laboratory abnormality of clinical concern has been identified. Investigators are also provided guidance on appropriate management of potential risks such as hypoglycemia (Section 6.4.9.1) urinary tract and genital infections (Section 6.4.9.2) hyponatraemia (Appendix E), decreased renal function (Appendix F) and increased liver function tests (Appendix G).

### **Potential benefits to subjects**

All subjects will receive active antihyperglycemic therapy including dapagliflozin monotherapy; both Japan phase II and global studies that have established the effect, but it needs to be confirmed in Japan phase III studies. In this study, the doses of dapagliflozin (5 mg and 10 mg) were chosen to provide efficacy in reducing hyperglycemia while mitigating the potential for AEs, based on previous clinical experience. In addition, dapagliflozin is expected to help decrease body weight (or prevent weight gain) as well as help lower blood pressure (BP) especially in subjects with elevated baseline BP. Subjects are also expected to receive some



benefit in the form of increased medical care/attention when participating in study procedures, which includes at least 15 clinic visits (17 clinical visits for subjects who need the washout period in the monotherapy group) with at least 14 physical examinations (16 physical examinations for subjects who need the washout period in the monotherapy group) over the 61-week study (the 67-week study for subjects who need the washout period in the monotherapy group).

### **Informed consent and alternatives to participation**

All prospective participants will be fully informed of the possible risks and benefits associated with this study, and their consent will be received prior to performing any study-related activity. Should a prospective participant elect to not participate in the study or to withdraw from the study, other medications are available to treat their diabetes, and the subject will not be disadvantaged in any way.

### **Conclusion**

Considering the pre-clinical and clinical experience with dapagliflozin and the precautions included in the study protocol, participation in this study presents a minimal and thus acceptable risk to subjects who meet the inclusion/exclusion criteria and consent to take part in the study.

## **2. STUDY OBJECTIVES**

### **2.1 Primary objective**

To evaluate the safety and tolerability of long term treatment up to 52 weeks with the dosing regimen of dapagliflozin, where it starts with 5 mg and titrates up to 10 mg depending on subjects' condition of glycemic control, for total subjects and for each subgroup of monotherapy or combination therapies with any anti-diabetic drugs including SU, rapid-acting insulin secretagogue (glinides), metformin,  $\alpha$ -glucosidase inhibitors (AGI), TZD, DPP-4 and GLP-1.

### **2.2 Secondary objectives**

To evaluate the efficacy of the dosing regimen of dapagliflozin for up to 52 weeks for each subgroup of monotherapy or combination therapy with SU, glinides, metformin, AGI, TZD, DPP4 or GLP-1 with regard to glycemic, weight and BP parameters.

### **2.3 Safety objective**

See Section [2.1](#).

## 2.4 Exploratory objectives – Not applicable

### 3. STUDY PLAN AND PROCEDURES

This Clinical Study Protocol has been subject to a peer review according to AstraZeneca standard procedures.

#### 3.1 Overall study design and flow chart

This is a long term, single arm, open label study to evaluate the safety and efficacy of dapagliflozin as monotherapy or in combination therapy. The study consists of only single arm without any comparator. However, the arm consists of 8 subgroups including monotherapy and combination therapies with SU, glinides, metformin, AGI, TZD, DPP-4 and GLP-1. The summary of each subgroup is outlined in Table 1. A total of 700 Japanese subjects with T2DM who have inadequate glycemic control on diet and exercise or on concomitant anti-diabetic treatment will be included in this study.

Either gender needs to be 40% or higher of total number of treated subjects for each sub-group. Only in the monotherapy group, the proportion of subjects with HbA1c  $\geq 6.5\%$  but  $\leq 7\%$  at Visit 5 needs to be at most approximately 25%.

**Table 1 Summary of subgroups**

Sub-group	Number of subjects	Background therapy			Inclusion criteria of HbA1c
		Drug	Dose before treatment started	Dose period before treatment started	
<b>Monotherapy</b>	240	-	-	-	6.5-10% at Visit 1 and 5 <sup>a</sup>
<b>Combination Therapy</b>					
SU	120	Glymeperide	Fixed dose within approved dose range in Japan	8 weeks	6.5-10% at Visit 1 and 5
Glinide	50	Any glinide			
Metformin	60	Any metformin			
AGI	60	Any AGI			
TZD	60	Pioglitazone		12 weeks	
DPP-4	60	Sitagliptin		8 weeks	
GLP-1	50	Liraglutide			

a Subjects who received ongoing medical treatment for diabetes within 6 weeks of enrolment, except for TZD (single oral anti-diabetic agent or two agents with less than half of the approved maximal dose for each agent) (see Inclusion criteria No.4) are required to have HbA1c  $\leq 8\%$  at Visit 1.


### 3.2 Monotherapy subgroup

For this subgroup, the study consists of a 2-week screening period, a 4-week lead-in period, a 52-week treatment period and a 3-week follow-up period. A 6-week wash-out period before a 4-week lead-in period is required in subjects who received ongoing medical treatment for diabetes within 6 weeks of enrolment, except for TZD (single oral anti-diabetic agent or two agents with less than half of the approved maximal dose for each agent) (see Inclusion criteria No.4).

Subjects who have inadequate glycemic control (HbA1c  $\geq 6.5\%$  but  $\leq 10\%$ ) on diet and exercise alone at Visit 5 (1 week before the start of the treatment period) are eligible to entry into the treatment period.

The study flow chart of monotherapy is shown in Figure 1.

**Figure 1 Study flow chart (monotherapy)**



E: Enrolment, T: Start of treatment

\* Wash-out period is applicable only for subjects who received medical treatment for diabetes up until enrolment. Drug-naïve subjects skip the period and directly proceed to the lead-in period.

\*\* Visit numbers correspond to those in subgroups of combination therapies. Visit 7 (Week 1) is missing here.

#### 3.2.1 Screening Period (2 weeks) : Visit 1-2

Subjects who are treatment-naïve (see Inclusion criteria No.4) are required to have HbA1c  $\geq 6.5\%$  but  $\leq 10\%$  and meet all inclusion and exclusion criteria at Visit1.

Subjects who received ongoing medical treatment for diabetes within 6 weeks of enrolment, except for TZD (single oral anti-diabetic agent or two agents with less than half of the approved maximal dose for each agent) (see Inclusion criteria No.4) are required to have HbA1c  $\leq 8\%$  at Visit1 and meet all inclusion and exclusion criteria. Subjects treated with TZD within 6 months prior to enrolment are not eligible for this study.

If the subjects have FPG > 240 mg/dL at Visit 1, the subjects are withdrawn from the study.

#### 3.2.2 Wash-out period (6 weeks) : Visit 2-4

Subjects, who are treatment-naïve or previously treated with anti-diabetic agents but have not been treated within 6 weeks of enrolment (see Inclusion criteria No.4), can skip the wash-out period and proceed to lead-in period.

Subjects who received ongoing medical treatment for diabetes within 6 weeks of enrolment, except for TZD (single oral anti-diabetic agent or two agents with less than half of the approved maximal dose for each agent) (see Inclusion criteria No.4) should have a 6-week washout period before a lead-in period (10-week washout before randomisation).

Washout means to stop taking oral anti-diabetic agent(s). Subjects will receive dietary counselling according to nutritional recommendations consistent with Japan Society of Diabetes or similar guidelines. Also, subjects will be given a blood glucose meter and diary, and be instructed on its use by site personnel.

If the subjects have FPG >240 mg/dL at Visit 2 or 3, the subjects are withdrawn from the study.

### **3.2.3 Lead-in Period (4 weeks) :Visit 4-6**

Eligible subjects are enrolled in a 4-week lead-in period while diet and exercise instruction will be provided.

Subjects will receive dietary counselling from the investigators according to nutritional recommendations consistent with Japan Society of Diabetes or similar guidelines. Subjects, who are treatment-naïve or previously treated with anti-diabetic agents but have not been treated within 6 weeks of enrolment (see Inclusion criteria No.4) (skipped wash-out period) will be given a blood glucose meter and diary, and be instructed on its use by site personnel.

If the subjects have FPG >240 mg/dL at Visit 4 or 5, the subjects are withdrawn from the study.

Subjects who have inadequate glycemic control (HbA1c  $\geq 6.5\%$  but  $\leq 10\%$ ) on diet and exercise alone at Visit 5 (1 week before the start of the treatment period) and who meet the eligibility criteria are eligible to entry into the treatment period at Visit 6.

### **3.2.4 Treatment Period (52 weeks):Visit 6-17**

Either gender needs to be 40% or higher of total number of treated subjects for the monotherapy sub-group. Also, the proportion of subjects with HbA1c  $\geq 6.5\%$  but  $\leq 7\%$  at Visit 5 needs to be at most approximately 25%.

Dapagliflozin starts with 5 mg at Visit 6. Up to Visit 10 (Week 12), the subject is treated with 5 mg of dapagliflozin.

At or after Visit 10 (Week 12), if the subject is under the conditions of HbA1c  $>7.5\%$  and there are no safety concerns, **at first**, the dose of dapagliflozin should be up-titrated to 10 mg from the next visit onward. After the dose up-titrated to 10 mg of dapagliflozin is administered for 8 weeks, the subject with inadequate glycemic control criteria outlined in [Table 2](#) should be considered for the rescue treatment by the investigator (See Section [3.2.4.1](#)).

Down titration to 5 mg of dapagliflozin is not allowed after up-titration to 10 mg of dapagliflozin. If the investigators judges the need of down titration to 5 mg of dapagliflozin, the subject must be withdrawn from the study.

### 3.2.4.1 Rescue Treatment

After the dose up-titrated to 10 mg of dapagliflozin is administered for 8 weeks, if HbA1c is still >8% in subjects from Visit 13 (Week 24) to Visit 17 (Week 52), the rescue treatment should be considered at the next visit, in principle (Table 2). An extra visit can be scheduled to initiate the rescue treatment, if the investigators judged it is needed as soon as possible. Thus, only one additional anti-diabetic drug will be added gradually up to the approved maximum doses at the discretion of investigators following local regulation and treatment guideline in Japan. If insulin is added, investigators should increase the dose with fully careful attention to avoid hypoglycemia caused by rapid increase.

Rescued subjects with central laboratory HbA1c values greater than 8% despite a maximum tolerated dose of rescue therapy for 12 weeks will be discontinued from the study and referred for additional anti-diabetic therapy (See Section 5.8).

**Table 2** Criteria for initiation of rescue therapy during the treatment period

Period	Central Laboratory HbA1c
From Visit 13 (Week 24) to Visit 17 (Week 52)	HbA1c >8%

During Visit 11 (Week 16) to Visit 13 (Week 24), if the results of self-monitored FPG show more than 240 mg/dL, the subject should repeat the FPG measurement on the same day. See Section 6.4.9.1, if the results of self-monitored FPG show more than 240 mg/dL consecutively.

### 3.2.5 Follow-up period (3 weeks) : Visit 17-18

Subjects will stop taking investigational products at Visit 17 (Week 52) and will be re-evaluated 3 weeks later for AEs and laboratory parameters. Subjects who discontinue study medication during the treatment period will also be re-evaluated 3 weeks later for AEs and laboratory parameters. Open-label use of anti-diabetic agents are permitted during the follow-up period.


### 3.3 Combination therapy subgroup

This study consists of a 2-week screening period, a 4-week lead-in period, a 52-week treatment period and a 3-week follow-up period.

Subjects who have inadequate glycemic control despite the treatment with basal anti-diabetic drug and required to have HbA1c  $\geq 6.5\%$  and  $\leq 10\%$  at Visit 1 and meet all inclusion and exclusion criteria are eligible to entry into the treatment period.

The study flow chart of combination Therapy is shown in [Figure 2](#).

**Figure 2 Study flow chart (combination therapies)**



E: Enrolment, T: Start of treatment

\* Required basal treatment period at stable doses before start of dapagliflozin treatment:  
At least 8 weeks; subgroups of SU, AGI, metformin, glinides, DPP-4 and GLP-1  
At least 12 weeks; subgroup of TZD

\*\* Visit 7 (Week 1) is applicable exclusively for the subgroup of combination therapy with GLP-1

\*\*\* Visit numbers correspond to those in monotherapy subgroup. Visit 2 and 3 are missing here

#### 3.3.1 Screening Period (2 weeks) : Visit 1 and 4

The subject has already been treated with fixed dose of basal anti-diabetic drug with a minimal 8 weeks or more on treatment within approved dose ranges in Japan before start of dapagliflozin. Twelve weeks treatment is required only in TZD subgroup.

Subjects must have inadequate glycemic control despite the treatment with basal anti-diabetic drug and required to have HbA1c  $\geq 6.5\%$  and  $\leq 10\%$  at Visit 1 and meet all inclusion and exclusion criteria.

The doses of basal anti-diabetic drug should not be changed during this period.

If the subjects have FPG  $>240$  mg/dL at Visit 1, the subjects are withdrawn from the study.

#### 3.3.2 Lead-in Period (4 weeks) : Visit 4-6

Eligible subjects are enrolled in a 4-week lead-in period while diet and exercise instruction will be provided.

Subjects will receive dietary counselling from the investigators according to nutritional recommendations consistent with Japan Society of Diabetes or similar guidelines. Also, subjects will be given a blood glucose meter and diary, and be instructed on its use by site personnel.

The doses of basal anti-diabetic drug should not be changed during this period.

If the subjects have FPG >240 mg/dL at Visit 4 or 5, the subjects are withdrawn from the study.

### **3.3.3 Treatment Period (52 weeks) : Visit 6-17**

Either gender needs to be 40% or higher of total number of treated subjects for each subgroup.

Dapagliflozin starts with 5 mg at Visit 6. Up to Visit 10 (Week 12), the subject is treated with 5 mg of dapagliflozin and with fixed dose of basal anti-diabetic drug. Subjects treated with GLP-1 will visit the study site at Visit 7 (Week 1).

At or after Visit 10 (Week 12), if the subject is under the conditions of HbA1c >7.5% and there are no safety concerns, **at first**, the dose of dapagliflozin should be up-titrated to 10 mg from the next visit onward. After the dose up-titrated to 10 mg of dapagliflozin is administered for 8 weeks, the subject with inadequate glycemic control criteria outlined in [Table 3](#) should be considered for the rescue treatment by the investigator (See Section [3.3.3.2](#)).

Down titration to 5 mg of dapagliflozin is not allowed after up-titration to 10 mg of dapagliflozin. If the investigator judges the need of down titration of dapagliflozin, the subject must be withdrawn from the study.

In addition, dose reduction of basal anti-diabetic drug (except for SU, See Section [3.3.3.1](#)) is not allowed during the treatment period at any condition. If the investigators judge the need of dose reduction of basal anti-diabetic drug (except for SU), the subject must be withdrawn from the study.

#### **3.3.3.1 Criteria for reduction of basal SU**

Reduction of SU dose shall be ordered by the investigators if the subject reports two or more readings of plasma glucose value  $\leq 70$  mg/dL ( $\leq 3.8$  mmol/L) at any visit after Visit 6.

In addition, the investigator may consider an SU dose reduction in subjects who report symptoms suggestive of hypoglycemia without confirmatory glucose measurements if, in the opinion of the investigators, the subject is at increased risk of further hypoglycemic episodes without such SU background dose adjustments.

#### **3.3.3.2 Rescue Treatment**

After the dose up-titrated to 10 mg of dapagliflozin is administered for 8 weeks, if HbA1c is still >8% in subjects from Visit 13 (Week 24) to Visit 17 (Week 52), the rescue treatment should be considered at the next visit, in principle ([Table 3](#)). An extra visit may be scheduled to initiate the rescue treatment, at the discretion of investigators. In principle, doses of the basal anti-diabetic drug will be gradually increased up to the approved maximum doses, and then if necessary, only

one additional anti-diabetic drug will be added gradually up to the approved maximum doses at the discretion of investigators following local regulation and treatment guideline in Japan. However, if the investigators have concern about increase of the doses of the original basal anti-diabetic drug in terms of safety, before reaching the maximum doses, only one additional anti-diabetic drug can be added first at the discretion of the investigators.

If insulin is added, investigators should increase the dose with fully careful attention to avoid hypoglycemia caused by rapid increase. Rescued subjects with central laboratory HbA1c values greater than 8% despite a maximum tolerated dose of rescue therapy for 12 weeks will be discontinued from the study and referred for additional anti-diabetic therapy (See Section 5.8).

**Table 3**                      **Criteria for initiation of rescue therapy during the treatment period**

<b>Period</b>	<b>Central Laboratory HbA1c</b>
From Visit 13 (Week 24) to Visit 17 (Week 52)	HbA1c >8.0%

During Visit 11 (Week 16) to Visit 13 (Week 24), if the results of self-monitored FPG show more than 240 mg/dL, the subject should repeat the FPG measurement on the same day. See Section 6.4.9.1, if the results of self-monitored FPG show more than 240 mg/dL consecutively.

### **3.3.4 Follow-up period (3 weeks) : Visit 17-18**

Subjects will stop taking investigational products at Visit 17 (Week 52) and will be re-evaluated 3 weeks later for AEs and laboratory parameters. Subjects who discontinue study medication during the treatment period will also be re-evaluated 3 weeks later for AEs and laboratory parameters. Open-label use of anti-diabetic agents are permitted during the follow-up period.



**Table 4 Study plan for subgroup of monotherapy**

Procedure	Screening Period	Washout Period <sup>a</sup>		Lead-in Period		Treatment Period										Follow up period		
	Wk -12 or Wk -6	Wk -10	Wk -7	Wk -4	Wk -1	Wk 0	Wk 4 <sup>b</sup>	Wk 8 <sup>b</sup>	Wk 12 <sup>b</sup>	Wk 16 <sup>b</sup>	Wk 20 <sup>b</sup>	Wk 24 <sup>b</sup>	Wk 32 <sup>b</sup>	Wk 40 <sup>b</sup>	Wk 48 <sup>b</sup>	Wk 52	Wk 55 <sup>b</sup>	
Visit	1	2	3	4	5	6	8	9	10	11	12	13	14	15	16	17	18	
Visit window (days)	+7	±3	±3	±3	±3	0	±3	±3	±3	±3	±3	±3	±3	±7	±7	±7	±7	±3
<b>Eligibility Assessments</b>																		
Informed Consent	X																	
Review Inclusion/Exclusion Criteria	X	X	X	X	X	X												
Complete Medical History	X																	
Complete Physical Examination	X					X						X				X		
Brief Physical Examination		X	X	X			X	X	X	X	X		X	X	X			X
Vital Signs, Body Weight	X			X		X	X	X	X	X	X	X	X	X	X	X	X	X
Height	X																	
Orthostatic BP						X	X	X				X				X	X	
Body Mass Index (BMI)	X					X						X				X	X	
Waist Circumference	X					X			X			X				X		
Electrocardiogram (ECG)	X					X						X				X		
Concomitant medications	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X
<b>General</b>																		
Dietary/exercise instruction/review		X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X
Dispense Glucose Meter, diary and provide instructions		X	X	X	X	X	X	X	X	X	X	X	X	X	X			

**Table 4 Study plan for subgroup of monotherapy**

Procedure	Screening Period	Washout Period <sup>a</sup>		Lead-in Period		Treatment Period										Follow up period		
	Wk -12 or Wk -6	Wk -10	Wk -7	Wk -4	Wk -1	Wk 0	Wk 4 <sup>b</sup>	Wk 8 <sup>b</sup>	Wk 12 <sup>b</sup>	Wk 16 <sup>b</sup>	Wk 20 <sup>b</sup>	Wk 24 <sup>b</sup>	Wk 32 <sup>b</sup>	Wk 40 <sup>b</sup>	Wk 48 <sup>b</sup>	Wk 52	Wk 55 <sup>b</sup>	
Visit	1	2	3	4	5	6	8	9	10	11	12	13	14	15	16	17	18	
Visit window (days)	+7	±3	±3	±3	±3	0	±3	±3	±3	±3	±3	±3	±3	±7	±7	±7	±7	±3
<b>Safety Assessments</b>																		
AE Assessment						X	X	X	X	X	X	X	X	X	X	X	X	X
SAE Assessment	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X
Hypoglycemic Events Assessment						X	X	X	X	X	X	X	X	X	X	X	X	X
<b>Laboratory Tests</b>																		
Pregnancy Test (urine) WOCBP only	X			X		X	X	X	X	X	X	X	X	X	X	X	X	X
Standard Safety Laboratory Panel (Blood and urine)	X			X		X	X	X	X	X	X	X	X	X	X	X	X	X
HbA1c	X				X	X	X	X	X	X	X	X	X	X	X	X	X	X
FPG	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X
Fasting Insulin, C-peptide	X					X		X	X	X	X	X	X	X	X	X		
Spot urine collection for glucose, sodium and creatinine	X					X			X	X	X	X	X	X	X	X		X
Fasting serum lipids: TC, LDL-C, HDL-C, TG and FFA						X			X			X		X		X		
Hepatitis Screen Panel <sup>c</sup> , TSH and PT/PTT	X																	
Cystatine C						X						X				X		
Parathormone (PTH)						X						X				X	X	

**Table 4 Study plan for subgroup of monotherapy**

Procedure	Screening Period	Washout Period <sup>a</sup>		Lead-in Period		Treatment Period										Follow up period		
	Wk -12 or Wk -6	Wk -10	Wk -7	Wk -4	Wk -1	Wk 0	Wk 4 <sup>b</sup>	Wk 8 <sup>b</sup>	Wk 12 <sup>b</sup>	Wk 16 <sup>b</sup>	Wk 20 <sup>b</sup>	Wk 24 <sup>b</sup>	Wk 32 <sup>b</sup>	Wk 40 <sup>b</sup>	Wk 48 <sup>b</sup>	Wk 52	Wk 55 <sup>b</sup>	
Visit	1	2	3	4	5	6	8	9	10	11	12	13	14	15	16	17	18	
Visit window (days)	+7	±3	±3	±3	±3	0	±3	±3	±3	±3	±3	±3	±3	±7	±7	±7	±7	±3
<b>Clinical Drug Supplies</b>																		
Dispense Study Medication						X	X	X	X	X	X	X	X	X	X	X	X	
Review Medication Compliance							X	X	X	X	X	X	X	X	X	X	X	

a Washout period is applicable only for subjects with ongoing treatment with anti-diabetic agents. For treatment-naïve subjects, screening period is Week -6.

b Subjects not completing the treatment period should complete the procedures described for Visit 17 within 7 days after discontinuation. These subjects should also complete the procedures described for a follow-up visit (ie, Visit 18) three weeks after discontinuation of investigational product.

c Includes anti-HAV (IgM, IgG), HBsAg, anti-HBc (IgG, IgM) and anti-HCV.

Note : Visit 7 (Week 1) is not applicable for subgroup of monotherapy.

WOCBP: women of child bearing potential, TC: total cholesterol, LDL-C: low-density lipoprotein cholesterol, HDL-C: high-density lipoprotein cholesterol , TG: triglyceride, FFA: free fatty acid, TSH: thyroid stimulating hormone, PT: prothrombin time, PTT: partial thromboplastin time,

**Table 5 Study plan for subgroups of combination therapy**

Procedure	Screening Period	Lead-in Period		Treatment Period												Follow up period	
	Wk -6	Wk -4	Wk -1	Wk 0	Wk 1	Wk 4 <sup>b</sup>	Wk 8 <sup>b</sup>	Wk 12 <sup>b</sup>	Wk 16 <sup>b</sup>	Wk 20 <sup>b</sup>	Wk 24 <sup>b</sup>	Wk 32 <sup>b</sup>	Wk 40 <sup>b</sup>	Wk 48 <sup>b</sup>	Wk 52	Wk 55 <sup>b</sup>	
Visit	1	4	5	6	7 <sup>a</sup>	8	9	10	11	12	13	14	15	16	17	18	
Visit Window (days)	+7	±3	±3	0	±3	±3	±3	±3	±3	±3	±3	±3	±7	±7	±7	±7	±3
<b>Eligibility Assessments</b>																	
Informed Consent	X																
Review Inclusion/Exclusion	X	X	X	X													
Complete Medical History	X																
Complete Physical Examination	X			X							X				X		
Brief Physical Examination		X			X	X	X	X	X	X		X	X	X			X
Vital Signs, Body Weight	X	X		X	X	X	X	X	X	X	X	X	X	X	X		X
Height	X																
Orthostatic BP				X	X	X	X					X			X	X	
BMI	X			X								X			X	X	
Waist Circumference	X			X				X				X			X		
ECG	X			X								X			X		
Concomitant medications	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X
<b>General</b>																	
Dietary/exercise instruction/review		X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X
Dispense Glucose Meter, diary and provide instructions		X	X	X		X	X	X	X	X	X	X	X	X			

**Table 5 Study plan for subgroups of combination therapy**

Procedure	Screening Period	Lead-in Period		Treatment Period												Follow up period	
	Wk -6	Wk -4	Wk -1	Wk 0	Wk 1	Wk 4 <sup>b</sup>	Wk 8 <sup>b</sup>	Wk 12 <sup>b</sup>	Wk 16 <sup>b</sup>	Wk 20 <sup>b</sup>	Wk 24 <sup>b</sup>	Wk 32 <sup>b</sup>	Wk 40 <sup>b</sup>	Wk 48 <sup>b</sup>	Wk 52	Wk 55 <sup>b</sup>	
Visit	1	4	5	6	7 <sup>a</sup>	8	9	10	11	12	13	14	15	16	17	18	
Visit Window (days)	+7	±3	±3	0	±3	±3	±3	±3	±3	±3	±3	±3	±7	±7	±7	±7	±3
<b>Safety Assessments</b>																	
AE Assessment				X	X	X	X	X	X	X	X	X	X	X	X	X	X
SAE Assessment	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X
Hypoglycemic Events Assessment				X	X	X	X	X	X	X	X	X	X	X	X	X	X
<b>Laboratory Tests</b>																	
Pregnancy Test (urine) WOCBP only	X	X		X		X	X	X	X	X	X	X	X	X	X	X	X
Standard Safety Laboratory Panel (Blood and urine)	X	X		X	X	X	X	X	X	X	X	X	X	X	X	X	X
HbA1c	X		X	X		X	X	X	X	X	X	X	X	X	X	X	X
FPG	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X
Fasting Insulin, C-peptide	X			X			X	X	X	X	X	X	X	X	X		
Spot urine collection for glucose, sodium and creatinine	X			X				X	X	X	X	X	X	X	X	X	X
Fasting serum lipids: TC, LDL-C, HDL-C, TG, and FFA				X				X				X		X		X	
Hepatitis Screen Panel <sup>c</sup> , TSH, and PT/PTT	X																
Cystatine C				X								X			X		
PTH				X								X			X	X	

**Table 5 Study plan for subgroups of combination therapy**

Procedure	Screening Period	Lead-in Period		Treatment Period												Follow up period	
	Wk -6	Wk -4	Wk -1	Wk 0	Wk 1	Wk 4 <sup>b</sup>	Wk 8 <sup>b</sup>	Wk 12 <sup>b</sup>	Wk 16 <sup>b</sup>	Wk 20 <sup>b</sup>	Wk 24 <sup>b</sup>	Wk 32 <sup>b</sup>	Wk 40 <sup>b</sup>	Wk 48 <sup>b</sup>	Wk 52	Wk 55 <sup>b</sup>	
Visit	1	4	5	6	7 <sup>a</sup>	8	9	10	11	12	13	14	15	16	17	18	
Visit Window (days)	+7	±3	±3	0	±3	±3	±3	±3	±3	±3	±3	±3	±7	±7	±7	±7	±3
<b>Clinical Drug Supplies</b>																	
Dispense Study Medication				X		X	X	X	X	X	X	X	X	X	X		
Review Medication Compliance						X	X	X	X	X	X	X	X	X	X	X	

a Visit 7 (Week 1) is applicable exclusively for subgroup of combination therapy with GLP-1.

b Subjects not completing the treatment period should complete the procedures described for Visit 17 within 7 days after discontinuation. These subjects should also complete the procedures described for a follow-up visit (ie, Visit 18) three weeks after discontinuation of investigational product.

c Includes anti-HAV (IgM, IgG), HBsAg, anti-HBc (IgG, IgM) and anti-HCV.

Note : Visit 2 and 3 are not applicable for subgroups of combination therapy.

### **3.4 Rationale for study design, doses and control groups**

#### **3.4.1 Study design and regulatory requirement**

The current study is designed to demonstrate the safety and tolerability of dapagliflozin as monotherapy or combination therapies in Japanese subjects of T2DM who have inadequate glycemic control.

##### **Monotherapy**

Subjects who received ongoing medical treatment for diabetes within 6 weeks of enrolment, except for TZD (single oral anti-diabetic agent or two agents with less than half of the approved maximal dose for each agent) will complete a 6-week washout period, and 4 weeks of lead-in for a total washout period of 10 weeks to eliminate effects of the prior therapy. The reason for 10 weeks of washout is as follows:

- (a) Adequate washout of all prior oral anti-diabetic agents and their pharmacodynamic effects
- (b) Stabilization of HbA1c after discontinuation of any oral anti-diabetic agents takes 8-12 weeks. 10 weeks of washout allows for an adequate stabilization of HbA1c after washout of oral anti-diabetic agents except for TZDs prior to treatment of subjects to dapagliflozin in this study.

The Follow-up period is set to examine the reversibility of potential AE and laboratory values after taking the last dose.

##### **Combination Therapy**

Considering the feature of each basal anti-diabetic drug, 8 weeks (12 weeks in TZD subgroup only) prior to treatment of dapagliflozin is required for an adequate stabilization of HbA1c.

The Follow-up period is set to examine the reversibility of potential AE and laboratory values after taking the last dose.

This study has standard design features for Long-term mono and combination therapy study (eg, multi-centre, open label) and incorporates the relevant features of the [Japanese Guideline on Clinical Evaluation methods on Oral Hypoglycemic Agents](#) which was issued 9<sup>th</sup> July 2010 with regard to duration of treatment, choice of study population, and choice of outcome variables.

#### **3.4.2 Study doses and control groups**

##### **Dapagliflozin**

Results of pre-clinical pharmacokinetic and toxicology studies support the safety of conducting a Phase III clinical development program for dapagliflozin. In Phase I clinical pharmacology studies (single ascending-dose study in healthy Japanese subjects and 2-week multiple ascending-dose study in Japanese subjects with T2DM), dapagliflozin was safe and

well tolerated with favourable pharmacokinetic and pharmacodynamic profile. A Japan Phase IIb study in subjects with T2DM demonstrated good glycemic efficacy and acceptable safety profile over a dose range from 1 mg to 10 mg. In addition, 5 mg and 10 mg of dapagliflozin showed clinically relevant effect on HbA1c (>0.5% placebo corrected reduction) over 12 weeks. Based on considerations of efficacy, pharmacodynamic, and safety data from the Phase I and II programs, doses of 5 mg and 10 mg of dapagliflozin have been chosen for the Phase III studies. The dose of dapagliflozin is expected to be increased during the long-term use in the clinical practice in Japan, so the titration dose regimen will be adopted.

### **Background therapy**

- Combination therapies with SU, glinides, metformin, AGI, DPP-4 or GLP-1  
The basal drugs are required to have been administered with stable doses within approved dose ranges for **8 weeks or more** before start of dapagliflozin treatment.
- Combination therapy with TZD  
Pioglitazone is required to have been administered with stable doses within approved dose ranges for **12 weeks or more** before start of dapagliflozin treatment.

### **3.4.3 Choice of outcome variables**

In the [Japanese Guideline on Clinical Evaluation methods on Oral Hypoglycemic Agents](#), dapagliflozin is expected to be long-term use in the clinical practice, therefore, the primary variable is to evaluate safety and tolerability of long term treatment up to 52 weeks with the dosing regimen of dapagliflozin as monotherapy or combination therapies with any anti-diabetic drugs approved and marketed.

### **3.4.4 Choice of study population**

#### **Age**

The prevalence of T2DM increases with age; it is therefore important to assess the safety of anti-diabetic agents in elderly subjects. In this study there is no upper age limit for subjects.

#### **HbA1c**

The HbA1c inclusion criterion was selected to include patients with a wide range of glycemic control. The lower limit of this range (ie, 6.5%) reflects the suggestion of the International Expert Committee, which was based on the results of DETECT2 study ([Gillett MJ, 2009](#)). The upper limit of this interval (ie, 10%) was chosen because insulin is generally the treatment of choice for patients with HbA1c values above this level ([ADA 2009](#)). To avoid potential bias in the subject characteristics which may be caused by excessive enrolment of subjects with low level of HbA1c, a limitation was added to the inclusion criterion, ie, enrolment of subjects with HbA1c  $\geq 6.5\%$  but  $\leq 7\%$  was limited to at most approximately 25% in the monotherapy subgroup. The measurement of HbA1c is National Glycohemoglobin Standardization Program (NGSP).



## **Pregnancy or breastfeeding**

Data from rat pre- and postnatal development and juvenile toxicity studies indicate that direct administration of dapagliflozin to weanling juvenile rats and indirect exposure during late pregnancy are each associated with increased incidence and/or severity of renal pelvic and tubular dilatations in progeny, although the long-term functional consequences of these effects are unknown. Dapagliflozin has not been tested in pregnant women and the risks to embryo, foetus, and infant are unknown. For this reason, women who are pregnant or breastfeeding are excluded and women of childbearing age are instructed to take precautions to avoid becoming pregnant during the study.

## **Other**

The purpose of the majority of the inclusion and exclusion criteria is to limit confounding factors that may complicate the interpretation of the study results (eg, corticosteroid-induced diabetes, haemoglobinopathies that would interfere with the HbA1c analyses) or to exclude subjects whose safety could be compromised by participation in the study.

## **4. SUBJECT SELECTION CRITERIA**

Investigator(s) should keep a record, the subject screening log, of subjects who entered pre-study screening.

Each subject should meet all of the inclusion criteria and none of the exclusion criteria for this study. Under no circumstances can there be exceptions to this rule.

### **4.1 Inclusion criteria**

For inclusion in the study subjects should fulfil the following criteria.

1. Provision of informed consent prior to any study specific procedures
2. Diagnosis of type 2 diabetes
3. Men or women age  $\geq 20$  years old (Either gender needs to be 40% or higher of total number of treated subjects for each subgroup)

Women not of childbearing potential (postmenopausal, and/or have undergone hysterectomy and/or bilateral oophorectomy or salpingectomy/tubal ligation) or women of childbearing potential who comply with the following:

- Use a highly effective method of birth control to avoid pregnancy throughout the study and for up to 4 weeks after the study
- Have a negative urine pregnancy test (minimum sensitivity 25 IU/L or equivalent units of human chorionic gonadotropin [HCG]) within 72 hours prior to the start of study medication and at each visit

Women will be defined as postmenopausal if last menstruation period was >1 year ago and serum follicle stimulating hormone (FSH) and luteinizing hormone (LH) are within the postmenopausal range, or if age >50 years and with last menstruation period >2 years ago.

4. Current condition at enrolment (Visit 1):

**For monotherapy subgroup**

- Drug naïve defined as:
  - Never received medical treatment for diabetes (insulin and/or other anti-diabetic agents [oral or injection])

**OR**

- Received medical treatment for diabetes for less than 30 days since diagnosis. In addition, during the 30-day period prior to screening did not receive oral anti-diabetic agents for more than 3 consecutive or more than 7 non-consecutive days. Subjects also should not have a history of insulin therapy within 2 weeks of screening (with the exception of insulin therapy during a hospitalization for other causes or use in gestational diabetes)

**OR**

- Previously received medical treatment for diabetes but have not been treated within 6 weeks of enrolment

**OR**

- Ongoing medical treatment for diabetes within 6 weeks of enrolment, except for TZD (single oral anti-diabetic agent or two agents with less than half of the approved maximal dose for each agent)

**For subgroups of combination therapies with SU, glinides, metformin, AGI, DPP-4 or GLP-1**

- Administration of each basal anti-diabetic drug at stable doses within approved dose ranges for 8 weeks or more before start of dapagliflozin treatment

**For subgroups of combination therapy with TZD**

- Administration of pioglitazone at stable doses within approved dose ranges for 12 weeks or more before start of dapagliflozin treatment

5. HbA1c:

**For monotherapy subgroup** (The proportion of subjects with HbA1c  $\geq 6.5\%$  but  $\leq 7\%$  at Visit 5 needs to be at most approximately 25%)

- At enrolment (Visit 1):
  - $\geq 6.5\%$  and  $\leq 10\%$  for subjects with drug naïve
  - $\leq 8\%$  for subjects with ongoing treatment
- At Week -1 (Visit 5) for treatment:
  - $\geq 6.5\%$  and  $\leq 10\%$  for all subjects

**For subgroups of combination therapies**

- At enrolment (Visit 1) for treatment:
  - $\geq 6.5\%$  and  $\leq 10\%$
- At Week -1 (Visit 5) for treatment:
  - $\geq 6.5\%$  and  $\leq 10\%$  for all subjects

## 4.2 Exclusion criteria

Subjects should not enter the study if any of the following exclusion criteria are fulfilled:

### Endocrine and metabolic disorders

1. Diagnosis of Type 1 diabetes mellitus, known diagnosis of maturity onset diabetes of the young (MODY) or secondary diabetes mellitus
2. History of diabetic ketoacidosis
3. Symptoms of poorly controlled diabetes including, but not limited to, marked polyuria, polydipsia, and/or greater than 10% weight loss during the 3 months prior to enrolment
4. FPG  $>240$  mg/dL (Visit 1, 2, 3, 4 and 5)
5. BMI  $\geq 45.0$  kg/m<sup>2</sup> at Visit 1
6. History of bariatric surgery (ie, any surgery to treat obesity; for example, gastric banding or procedures that involve bypassing or transposing sections of the small intestine). History of liposuction is allowed.
7. Diabetes insipidus

8. Thyroid-stimulating hormone (TSH) values outside normal range at Visit 1. An abnormal TSH value needs to be followed up with a free T4 test. Subjects with abnormal free T4 values will be excluded. Subjects with an elevated TSH and normal free T4 will be allowed if referred to their family doctor for thyroid hormone replacement therapy prior to treatment started

### **Kidney disorders**

9. Estimated glomerular filtration rate (eGFR): <45 mL/min (calculated by Japanese guideline formula) or a measured serum creatinine (SCr) value of >1.5 mg/dL (133 µmol/L) for male subjects and >1.4 mg/dL (124 µmol/L) for female subjects at Visit 1.

For subgroup of combination therapy with metformin: eGFR: <60 mL/min (calculated by Japanese guideline formula) or a measured SCr value of >1.5 mg/dL (133 µmol/L) for male subjects and >1.4 mg/dL (124 µmol/L) for female subjects at Visit 1.

10. Urine albumin: creatinine ratio (UACR) >1800 mg/g (>203.4 mg/mmol) at Visit 1
11. History of unstable or rapidly progressing kidney disease
12. Familial renal glucosuria. This condition is diagnosed as glucosuria (glucose >1.0 mmol/L urine) in the presence of normoglycemia in a subject without the diagnosis of diabetes mellitus

### **Hepatic disorders**

13. Severe hepatic insufficiency and/or significant abnormal liver function defined as aspartate aminotransferase (AST) >3X upper limit of normal (ULN) and/or alanine aminotransferase (ALT) >3X ULN at Visit 1
14. Total bilirubin (TB) >2.0 mg/dL (>34.2 µmol/L) at Visit 1
15. Positive serologic evidence of current infectious liver disease including Hepatitis A viral antibody IgM, Hepatitis B surface antigen and Hepatitis C virus antibody at Visit 1
16. History of drug-induced liver enzyme elevation
17. History of severe hepatobiliary disease or hepatotoxicity with any medication

### **Cardiovascular disorders**

18. Congestive heart failure defined as New York Heart Association (NYHA) class IV, unstable or acute congestive heart failure. Note: eligible subjects with congestive heart failure, especially those who are on diuretic therapy, should have careful monitoring of their volume status throughout the study

19. Significant cardiovascular history within the past 2 months prior to enrolment, defined as: myocardial infarction, unstable angina pectoris, transient ischemic attack, unstable or previously undiagnosed arrhythmia, cardiac surgery or revascularization (coronary angioplasty or bypass grafts), or cerebrovascular accident. In addition, subjects who have unstable cardiovascular disease at enrolment in the judgment of the investigators are excluded from the study.
20. Systolic blood pressure (SBP)  $\geq 160$  mmHg and/or diastolic blood pressure (DBP)  $\geq 100$  mmHg at Visit 1

#### **Hematologic/oncologic disorders/conditions**

21. Haemoglobin  $< 10$  g/dL ( $< 100$  g/L) for men; haemoglobin  $< 9$  g/dL ( $< 90$  g/L) for women at Visit 1
22. History of chronic haemolytic anaemia or haemoglobinopathies (for example, sickle cell anaemia, thalassemia, sideroblastic anaemia)
23. Iron deficiency anemia with iron therapy started in the past 12 weeks prior to enrolment, or a recent diagnosis of iron deficiency anemia that requires therapeutic management within the next 6 months in the judgement of the investigators
24. Donation or transfusion of blood, plasma, or platelets within the past 3 months prior to enrolment
25. History of malignancy within the last 5 years prior to enrolment, excluding successful treatment of basal or squamous cell skin carcinoma

#### **Infectious disease/immunologic disorders**

26. Known immunocompromised status, including subjects who have undergone organ transplantation

#### **Musculoskeletal disorders**

27. Creatinine Kinase (CK)  $> 3X$  ULN at Visit 1
28. History of drug-induced myopathy or drug-induced CK elevation

#### **Reproductive status**

29. Pregnant or breastfeeding subjects

#### **Prohibited medications**

30. Treatment with TZD within 6 month prior to enrolment (applicable for monotherapy group)

31. Use of weight loss medication, including but not limited to mazindol, sibutramine, phentermine, orlistat, rimonabant, benzphetamine, diethylpropion, methamphetamine, and/or phendimetrazine, within 30 days prior to enrolment
32. Treatment with systemic glucocorticoids equivalent to oral prednisolone  $\geq 10$  mg (betametasone  $\geq 1.2$  mg, dexamethasone  $\geq 1.5$  mg, hydrocortisone  $\geq 40$  mg) per day within 30 days prior to enrolment; single systemic doses, topical or inhaled corticosteroids are allowed
33. Treatment with unstable doses of teriparatide, bisphosphonates and/or calcitonin within 30 days prior to enrolment
34. Treatment for Human Immunodeficiency Virus (HIV) and/or use of antiviral drugs (delavirdine, indinavir, nelfinavir, ritonavir, saquinavir)

**Other**

35. Intolerance, contraindication or potential allergy or hypersensitivity to dapagliflozin
36. Any clinically significant abnormality identified on physical examination, ECG or laboratory tests, which in the judgement of the investigators would compromise the subject's safety or successful participation in the clinical study
37. Subjects who, in the judgement of the investigators, may be at risk for dehydration
38. Acute or chronic metabolic acidosis
39. History of alcohol abuse or illegal drug use within the past 12 months prior to enrolment
40. Involvement in the planning and conduct of the study (applies to both AstraZeneca and Bristol-Myers Squibb staff or staff at the study centre)
41. Previous enrolment or randomisation to treatment in the present study
42. Previous participation in a clinical study with dapagliflozin (BMS-512148) in which the subject received at least one dose of study medication except for placebo
43. Participation in another clinical study during the last 1 month prior to enrolment

Procedures for withdrawal of incorrectly enrolled subjects see Section 5.3.

## **5. STUDY CONDUCT**

### **5.1 Restrictions during the study**

- Subjects must be in a fasting state at least 10 hours prior to study visit (however, drinking water is allowed). Permitted medications may be taken with water only.
- Subjects must abstain from tobacco for 12 hours prior to all study visits.
- Subjects must make every attempt to adhere to the dietary and physical activity changes and goals as discussed with the Investigator(s).
- Women of child-bearing potential must immediately contact the Investigator if they suspect they might be pregnant and if they have changed, or plan to change their birth control method.
- The subjects should not take investigational products and basal anti-diabetic drug on the morning of the clinic visit.

If a subject arrives for a visit without having followed the above instructions, the entire visit should be rescheduled (within the allowed time-window, if possible).

As up to approximately 246 mL of blood will be drawn from each subject during the entire duration of the clinical study, subjects should be instructed to abstain from donating any blood during the clinical study and for 3 months following their last study visit.

Prohibited and restricted concomitant medications are listed in Section 5.6.

### **5.2 Subject enrolment**

The Principal Investigator will:

1. Obtain signed informed consent from the potential subject before any study specific procedures are performed.
2. Perform enrolment call in Interactive Web Response System (IWRS).
3. Assign potential subject a unique enrolment number “EXXXYYYY”, which is composed of 4 digits (XXXX) of centre number and 3 digits (YYY) of consecutive number in order of enrolment registration at each study site.
4. Determine subject eligibility. See Sections 4.1 and 4.2.
5. Perform call for start of treatment of dapagliflozin at Visit 6 in IWRS.

6. Assign eligible subjects with a unique subject number through IWRS. (Subject number will be assigned strictly sequentially as subjects become eligible for treatment.)

Routines for this will be described in the IWRS user manual that will be provided to each centre.

If a subject withdraws from participation in the study, then his/her enrolment number/subject number cannot be reused.

### **5.3 Procedures for handling subjects incorrectly enrolled or treated**

**Subjects who fail to meet the inclusion/exclusion criteria** should not, under any circumstances, be enrolled or treated. There can be no exceptions to this rule.

Where subjects that do not meet the selection criteria are treated in error or incorrectly started on treatment, or where subjects subsequently fail to meet the study criteria post initiation, a discussion should occur between the AstraZeneca Study Delivery Team Physician and the Investigator regarding whether to continue or discontinue the subject from treatment.

The AstraZeneca Study Delivery Team Physician is to ensure all such decisions are appropriately documented. In situations where an agreement cannot be reached, the subject should have their study therapy stopped.

### **5.4 Blinding and procedures for unblinding the study – Not applicable**

### **5.5 Treatments**

#### **5.5.1 Identity of investigational product(s)**

Dapagliflozin (BMS-512148) tablets are packed into HDPE bottle and are supplied by AstraZeneca or their designee. For details of the identity of the investigational product see below.

**Table 6 Identity of investigational product(s)**

<b>Investigational product</b>	<b>Dosage form and strength</b>	<b>Manufacturer</b>
BMS-512148 tablet 5 mg	Contains 5 mg of dapagliflozin and Green, plain diamond shaped, film coated tablet	Bristol-Myers Squibb Company
BMS-512148 tablet 10 mg	Contains 10 mg of dapagliflozin and Green, plain diamond shaped, film coated tablet	Bristol-Myers Squibb Company



### 5.5.2 Doses and treatment regimens

Investigational drug should be taken once daily in the morning.

- Dapagliflozin 5 mg dose: Dapagliflozin 5 mg 1 tablet
- Dapagliflozin 10 mg dose: Dapagliflozin 10 mg 1 tablet

**Table 7 Drug dispensing scheme**

Visit ID	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16
Number of bottles to dispense dapagliflozin 5 mg or 10 mg	0	0	0	0	0	1	0	1	1	1	1	1	2	2	2	1

### 5.5.3 Additional study drug – Not applicable

### 5.5.4 Labelling

Labelling of the investigational product will be carried out by AstraZeneca or Contract Research Organisation (CRO) in accordance with current Good Manufacturing Practice (GMP). Labels will be prepared in accordance with GMP and local regulatory guidelines. The labels will fulfil GMP Annex 13 requirements for labelling and text will be translated into local language. Details of labelling and packaging of the study drug will be described in a separate document, 'Procedure of storage conditions for investigational product'.

### 5.5.5 Storage

All study drugs should be kept in a secure place under appropriate storage conditions. A description of the appropriate storage conditions is specified on the investigational product pack label.

## 5.6 Concomitant and post-study treatment(s)

Changes in concomitant medication should be avoided during study participation, with the exception of situations defined in this protocol, but medication, which is considered necessary for the subject's safety and well-being, may be given at the discretion of the investigators, who must decide if the subject should remain in study or need to be dismissed from study due to subject's safety or interference with study objectives.

The administration of all medication must be recorded in the appropriate sections of the electronic case report form (eCRF) with trade names, dosages, dates of starting and ending of medication and reason for therapy.

After having completed or discontinued the study, subjects will receive usual care and anti-diabetic agents according to the investigator's judgement and according to local medical practice.

### **Prohibited Therapies**

- Weight loss medication, including but not limited to mazindol, sibutramine, phentermine, orlistat, rimonabant, benzphetamine, diethylpropion, methamphetamine, and/or phendimetrazine
- Antiviral drugs (delavirdine, indinavir, nelfinavir, ritonavir, saquinavir)
- Treatment with systemic glucocorticoids equivalent to oral prednisolone  $\geq 10$  mg (betametasone  $\geq 1.2$  mg, dexamethasone  $\geq 1.5$  mg, hydrocortisone  $\geq 40$  mg) per day (two temporary periods of higher daily doses for no longer than 7 days each are allowed).
- Anti-diabetic agents except for the study medication, a basal anti-diabetic agents medication and rescue therapies (Anti-diabetic agents planned to be stopped for wash-out are allowed to be used until the beginning of wash-out period. After completion of the treatment period, alternative anti-diabetic treatment will be initiated according to standard medical practice.)

### **Restricted Therapies**

- Insulin use is permitted in the following cases:
  - For up to 14 days total during the study and up to 7 continuous days if subjects are unable to take oral medications (for example during a gastrointestinal illness)
  - For up to 14 days total during the study and up to 7 continuous days if there is a documented illness or infection that requires additional therapy for maintaining glycemic control
  - For up to 14 days total during the study and up to 7 continuous days if subjects have to temporarily stop investigational product and rescue therapy due to requirements of this clinical study protocol.
  - For up to 7 days during hospitalisation as long as the primary reason for hospitalisation is not management of the subject's glycemic control.
- Teriparatide, bisphosphonates and/or calcitonin are allowed if only the dose has not changed.

## **5.7 Treatment compliance**

The administration of all study drugs (including investigational products) should be recorded in the appropriate sections of the eCRF.

Subjects will be asked to return all unused investigational product and empty packages and bottles to the clinic at each visit. The subject will be asked about compliance at each study

visit; compliance will also be assessed based on returned tablet counts. Tablet counts will be recorded in the eCRF. Subjects judged to be non-compliant (defined as taking less than 80% or more than 120% of the prescribed dose of investigational product) may continue in the study, but should be counselled on the importance of taking their study medication as prescribed.

### **5.7.1 Accountability**

Study drug will not be distributed to the study site until the contract is concluded between the study site and AstraZeneca. The Investigational Product Storage Manager is responsible for managing the study drug from receipt by the institution until the return of all unused study drug to AstraZeneca. AstraZeneca will provide the study documents 'Procedures for drug accountability' and 'Procedures for drug storage' which describes the specific requirements. The investigator(s) is responsible for ensuring that the subject has returned all unused study drug.

## **5.8 Discontinuation of investigational product**

Subjects who discontinue study medication during the treatment period will be performed the procedures scheduled at Visit 17 (Week 52) and will be re-evaluated 3 weeks later for AE. Open-label use of anti-diabetic agents are permitted during the follow-up period.

### **5.8.1 Criteria for discontinuation from the study**

Subjects may be discontinued from study treatment and assessments at any time. Specific reasons for discontinuing a subject are:

#### **General discontinuation criteria:**

1. Voluntary discontinuation by the subject who is at any time free to discontinue his/her participation in the study, without prejudice to further treatment
2. Risk to subjects as judged by the investigator, AstraZeneca and/or Bristol-Myers Squibb
3. Severe non-compliance to protocol as judged by the investigator, AstraZeneca and/or Bristol-Myers Squibb
4. Incorrectly enrolled subjects (See Section 5.3)
5. Subject lost to follow-up defined by inability to reach the subject after 3 documented phone calls, faxes, or e-mails. All attempts at contact should be documented in the subject's medical record
6. AEs, ie, any clinical AE, laboratory abnormality or intercurrent illness which, in the opinion of the investigator, indicates that continued participation in the study is not in the best interest of the subject

**Study-specific discontinuation criteria:**

7. Use of (need for) any anti-diabetic agents, except for rescue treatment, other than investigational product and/or basal anti-diabetic agents.
8. Treatment with systemic glucocorticoids equivalent to oral prednisolone  $\geq 10$  mg/day (two temporary periods of higher daily doses for no longer than 7 days each are allowed)
9. Major and/or frequent hypoglycemic events; If more than one event of severe hypoglycemia, defined as symptomatic events requiring external assistance due to severe impairment in consciousness or behaviour, with a capillary or plasma glucose value  $< 3$  mmol/L ( $< 54$  mg/dL), and prompt recovery after glucose or glucagon administration occurs, or if the investigators decides that current treatment poses the subject at risk of either repeated hypoglycemic events or severe hypoglycemia.
10. Pregnancy confirmed by a positive pregnancy test or otherwise verified
11. Rescued subjects with central laboratory HbA1c values greater than 8.0% despite a maximum tolerated dose of rescue therapy for 12 weeks
12. Change in kidney function (Please see Appendix F for further guidance):
  - For subjects treated with metformin: For subjects with a central laboratory eGFR  $< 60$  mL/min or with an increase in SCr of  $\geq 0.5$  mg/dL ( $44.2$   $\mu\text{mol/L}$ ) above the baseline value (Visit 6), an increase in SCr of  $\geq 0.5$  mg/dL above the baseline value (Visit 6) confirmed by a repeated measurement within one week or eGFR  $< 60$  mL/min
  - For subjects not treated with metformin: Subject with following criteria will be scheduled for an additional visit within one week.
    - (a) eGFR  $< 45$  ml/min
    - (b) Subjects with baseline creatinine  $\geq 1.4$  mg/dL ( $\geq 123$   $\mu\text{mol/L}$ ) an absolute increase in SCr of  $\geq 1.0$  mg/dL ( $\geq 88$   $\mu\text{mol/L}$ ), or
    - (c) Subjects with baseline creatinine  $< 1.4$  mg/dL ( $< 123$   $\mu\text{mol/L}$ ) an absolute increase in SCr of  $\geq 0.5$  mg/dL ( $\geq 44$   $\mu\text{mol/L}$ ).

Subjects should be discontinued from study if the repeat laboratory tests meet SCr  $\geq 0.5$  mg/dL or eGFR  $< 45$  mL/min.
13. Subjects with a central laboratory CK  $> 10$ X ULN will be scheduled for an additional visit within 24 hours, but not exceeding 72 hours. Subjects should be discontinued from study if the repeat laboratory tests meet CK  $> 10$ X ULN.

14. Subjects with a central laboratory ALT and/or AST >3X ULN will be scheduled for a follow-up visit within 3 days following the receipt of the result. See Appendix G for further guidance. Subjects should be discontinued from study if the initial and repeat laboratory tests meet any of the following criteria:
- (a) ALT and/or AST are >3X ULN and TB >1.5X ULN
  - (b) ALT and/or AST are >5X ULN for  $\geq 14$  consecutive days, at any time after initial confirmatory results
  - (c) ALT and/or AST are >8X ULN.
15. Serum Sodium  $\leq 125$  mmol/L with or without symptoms (see Appendix E for further guidance)

### 5.8.2 Procedures for discontinuation of a subject from investigational product

A subject that discontinues will always be asked about the reason(s) for discontinuation and the presence of any AEs. The Principal Investigator/sub-investigator will perform the best possible observation(s), test(s) and evaluation(s) as well as give appropriate medication and all possible measures for the safety of the subject. They will also immediately inform AstraZeneca of the withdrawal. AEs will be followed up (See Sections 6.4.3 and 6.4.4); diary, glucometer and study drug should be returned by the subject.

#### Pre-treatment

A subject that discontinues will always be asked about the reason(s) for discontinuation and the presence of any AEs. If possible, they will be seen and assessed by an investigator(s). AEs will be followed up (See Sections 6.4.3 and 6.4.4); subject diaries, glucometer and study drug should be returned by the subject.

#### Treated subjects

Treated subjects who discontinue the study treatment before Week 52 should return and complete the procedures described for Visit 17 as soon as possible but at the latest 7 days after discontinuation. These subjects should also be scheduled for a follow-up visit (ie, procedures of Visit 18) three weeks after discontinuation of investigational product. In addition subjects who discontinue the study due to an AE including a laboratory abnormality should be followed by the investigator until the event has been resolved or stabilised.

After discontinuation of investigational product, alternative anti-diabetic treatment will be initiated according to the investigator's judgement and according to local medical practice.

Subjects with an increased CK >10X ULN will have their investigational product held and repeated CK test preferably within 24 hours, but not exceeding 72 hours. If repeated CK is still >10X ULN the subject should permanently discontinue all study medication and be withdrawn from the study (in which case an AE must be reported). Otherwise investigational product may be resumed unless otherwise contraindicated.

Subjects with increased liver function tests as defined in Section 5.8.1 under listing No.14 will be scheduled for a follow-up visit within 3 days following the receipt of the result. Subjects may remain on study medication until the confirmatory results are obtained. If repeat liver function tests still are increased as outlined in Appendix G the subject should permanently discontinue study medication and be withdrawn from the study (See Appendix G for further guidance).

## **5.9 Withdrawal from study**

Subjects are at any time free to withdraw from study (investigational product and assessments), without prejudice to further treatment (withdrawal of consent). Such subjects will always be asked about the reason(s) and the presence of any AEs. If possible, they will be seen and assessed by an investigator. AEs will be followed up (See Sections 6.4.3 and 6.4.4); diary, glucometer and all study drugs should be returned by the subject.

## **6. COLLECTION OF STUDY VARIABLES**

### **6.1 Recording of data**

The Rave Web Based Data Capture (WBDC) system will be used for data collection and query handling. The investigator will ensure that data are recorded on the eCRFs as specified in the study protocol and in accordance with the instructions provided.

The investigator ensures the accuracy, completeness and timeliness of the data recorded and of the provision of answers to data queries. The investigator will sign the completed eCRF. A copy of the completed eCRFs will be archived at the study site.

The Principal Investigator/sub-investigator will record data on the observations, tests and assessments specified in the protocol on the eCRFs provided by AstraZeneca. The eCRF will be accompanied with 'Instructions for the Investigator', which should be followed. These instructions provide guidance for the recording of study data in the eCRF including how to change data incorrectly recorded.

### **6.2 Data collection and enrolment**

For full details please see the study plan (Table 4 or Table 5).

#### **6.2.1 Follow-up procedures**

A follow-up (Visit 18) will be performed 3 weeks ( $\pm 3$  days) after the end of the treatment period, see Table 4 or Table 5 for further details.

## 6.3 Efficacy

### 6.3.1 Efficacy laboratory variables

The laboratory parameters that will be measured to assess efficacy are displayed in [Table 8](#) by visit.

**Table 8 Efficacy laboratory variables**

Visit	1	2 <sup>a</sup>	3 <sup>a</sup>	4	5	6	7 <sup>b</sup>	8	9	10	11	12	13	14	15	16	17	18
Study week	-12 or -6	-10	-7	-4	-1	0	1	4	8	12	16	20	24	32	40	48	52	55
HbA1c	X				X	X		X	X	X	X	X	X	X	X	X	X	X
FPG	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X
Insulin	X					X			X	X	X	X	X	X	X	X	X	X
C-peptide	X					X			X	X	X	X	X	X	X	X	X	X
TC						X				X			X		X		X	
LDL-C						X				X			X		X		X	
HDL-C						X				X			X		X		X	
TG						X				X			X		X		X	
FFA						X				X			X		X		X	

a Visit 2 and 3 are only applicable for subgroup of monotherapy.

b Visit 7 (Week 1) is only applicable for subgroup of GLP1 in combination therapy.

### 6.3.2 HbA1c

HbA1c is a well established measure of glycemic efficacy referred by the [Japanese Guideline on Clinical Evaluation methods on Oral Hypoglycemic Agents](#) and considered to be an acceptable secondary endpoint the assessment for the determination of glycemic efficacy. HbA1c will be analysed by a central laboratory according to the procedures described in the Laboratory Manual which will be distributed to each study site during site initiation. The measurement of HbA1c is NGSP.

### 6.3.3 FPG

FPG is a well established measure of glycemic efficacy and considered to be an acceptable secondary endpoint.

### 6.3.4 Weight and height

The subject's weight will be recorded in kilogram (kg) to one decimal place, on a fasting stomach with light clothing and no shoes. The subject's height will be recorded in centimetres, with no shoes. All readings should be recorded as accurately as possible and the same scale should be used for all assessments for a given subject.

### **6.3.5 BMI**

BMI is a calculated ratio between weight and height (weight / height<sup>2</sup>, where weight is measured in kg, and height in metres) and will be computed by AstraZeneca.

### **6.3.6 Blood pressure**

As BP is both an efficacy and safety variable in this study, measurement of BP is described in Sections [6.4.8.1](#) and [6.4.8.2](#).

### **6.3.7 Waist circumference**

The waist circumference should be measured in the standing position at the natural waist (smallest waist circumference). If there is no natural waist, the measurement should be made at the level of the umbilicus. Measurements should be made at the end of a normal inspiration.

## **6.4 Safety**

The Principal Investigator is responsible for ensuring that all staff involved in the study is familiar with the content of this section.

### **6.4.1 Definition of adverse events**

An AE is the development of an undesirable medical condition or the deterioration of a pre-existing medical condition following or during exposure to a pharmaceutical product, whether or not considered causally related to the product. An undesirable medical condition can be symptoms (eg, nausea, chest pain), signs (eg, tachycardia, enlarged liver) or the abnormal results of an investigation (eg, laboratory findings, electrocardiogram). In clinical studies, an AE can include an undesirable medical condition occurring at any time, including run-in or washout periods, even if no study treatment has been administered.

The term AE is used to include both serious and non-serious AEs.

For cases where it could be suspected that a tissue-derived medicine has been contaminated by a pathogen, information about any of the above conditions (including infection) should be collected.

### **6.4.2 Definitions of serious adverse event**

A SAE is an AE occurring during any study phase (ie, run-in, treatment, washout, follow-up), that fulfils one or more of the following criteria:

- Results in death
- Is immediately life-threatening
- Requires in-subject hospitalisation or prolongation of existing hospitalisation (including hospitalisation for tests related to AEs), except hospitalisation that has been planned before enrolment



- Results in persistent or significant disability or incapacity
- Is a congenital abnormality or birth defect
- Is an important medical event that may jeopardise the subject or may require medical intervention to prevent one of the outcomes listed above.
- Cancer
- Drug Dependency/ abuse
- Overdose (defined as the accidental or intentional ingestion of any dose of the investigational product that is considered both excessive and medically important)

For further guidance on the definition of a SAE, see Appendix B to the Clinical Study Protocol.

### **6.4.3 Recording of adverse events**

#### **Time period for collection of adverse events**

AEs will be collected throughout the treatment period (Visit 6 to 17) and including the follow-up period (Visit 18).

SAEs will be recorded from the time of informed consent is obtained until the end of the study (Visit 18).

#### **Follow-up of unresolved adverse events**

Any AEs that are unresolved at the end of the study (Visit 18) or at discontinuation visit in the study are followed up by the investigator for as long as medically indicated, but without further recording in the eCRF. AstraZeneca retains the right to request additional information for any subject with ongoing AE(s)/SAE(s) at the end of the study, if judged necessary.

#### **Variables**

The following variables will be collect for each AE;

- AE (verbatim)
- the date when the AE started and stopped
- maximum intensity
- whether the AE is serious or not
- investigator causality rating against the Investigational Product (yes or no)
- action taken with regard to investigational product

- outcome.

In addition, the following variables will be collected for SAEs:

- Date AE met criteria for serious AE
- Date Investigator became aware of serious AE
- AE is serious due to
- Date of hospitalisation
- Date of discharge
- Probable cause of death
- Date of death
- Autopsy performed
- Causality assessment in relation to Study procedure(s)
- Causality assessment in relation to Other medication
- Causality assessment in relation to Additional Study Drug
- Description of AE

Maximum intensity will be graded according to the following definitions:

- mild (awareness of sign or symptom, but easily tolerated)
- moderate (discomfort sufficient to cause interference with normal activities)
- severe (incapacitating, with inability to perform normal activities)
- very severe (debilitating, significantly incapacitates subject despite symptomatic therapy)

It is important to distinguish between serious and severe AEs. Severity is a measure of intensity whereas seriousness is defined by the criteria in Section 6.4.2. An AE of severe intensity need not necessarily be considered serious. For example, nausea that persists for several hours may be considered severe nausea, but not a SAE. On the other hand, a stroke

that results in only a limited degree of disability may be considered a mild stroke but would be a SAE.

### **Causality collection**

The Investigator will assess causal relationship between Investigational Product and each AE, and answer 'yes' or 'no' to the question 'Do you consider that there is a reasonable possibility that the event may have been caused by the investigational product?'

For SAEs causal relationship will also be assessed for other medication and study procedures and additional study drug (rescue medication). Note that for SAEs that could be associated with any study procedure the causal relationship is implied as 'yes'.

A guide to the interpretation of the causality question is found in Appendix B to the Clinical Study Protocol.

### **Adverse Events based on signs and symptoms**

All AEs spontaneously reported by the subject or reported in response to the open question from the study personnel: 'Have you had any health problems since the previous visit/you were last asked?', or revealed by observation will be collected and recorded in the eCRF. When collecting AEs, the recording of diagnoses is preferred (when possible) to recording a list of signs and symptoms. However, if a diagnosis is known and there are other signs or symptoms that are not generally part of the diagnosis, the diagnosis and each sign or symptom will be recorded separately.

### **Adverse Events based on examinations and tests**

Deterioration as compared to baseline in protocol-mandated laboratory values, vital signs and other safety variables will only be reported as AEs if they are clinically significant, fulfil any of the SAE criteria or are the reason for discontinuation of treatment with the investigational product, or require the subject to receive specific corrective therapy. If a deterioration in a laboratory value/vital sign is associated with clinical signs and symptoms, the sign/symptom will be reported as an AE and the associated laboratory result/vital sign will be considered as additional information. Any new or aggravated clinically relevant abnormal medical finding at a physical examination or ECG evaluation as compared with the baseline assessment will be reported as an AE.

Clinically relevant deterioration in non-protocol-mandated measurements will be reported as AE(s).

Wherever possible the reporting investigator uses the clinical, rather than the laboratory term (eg, anaemia versus low haemoglobin value).

### **Hypoglycemic events**

A separate section in the eCRF will be used to document all reported episodes of hypoglycemia (See Section 6.4.9). Hypoglycemic episodes should only be reported on the AE

eCRF page if the event fulfils protocol criteria for a SAE (See Section 6.4.2). In this case, an SAE must be reported in addition to the hypoglycemia eCRF pages for hypoglycemia.

### **Follow-up of unresolved adverse events**

All AEs and SAEs, including those that are ongoing at the end of the study or at discontinuation, will be followed up until resolution or until the Investigator decides that no further follow-up is necessary. AstraZeneca retains the right to request additional information for any subject with ongoing AEs/SAEs at the end of the study, if judged necessary. The requirement to follow-up is not intended to delay database lock or production of the clinical study report. Both these activities should proceed as planned with ongoing AEs if necessary.

Any follow-up of ongoing SAEs after database lock will be reported to AstraZeneca, who will notify the appropriate Bristol-Myers Squibb Pharmacovigilance contact.

### **Cardiovascular events**

Cardiovascular events will be monitored in the study population and an independent cardiovascular adjudication committee will review events. Cardiovascular events will be analyzed in conjunction with cardiovascular events observed in other Phase II and Phase III dapagliflozin studies and reported separately.

#### **6.4.4 Reporting of serious adverse events**

Investigators and other study site personnel must inform appropriate AstraZeneca representatives via the WBDC system of any SAE that occurs in the course of the study within 1 calendar day (ie, immediately but no later than the end of the next business day) of when he or she becomes aware of it. Follow-up information on SAEs must also be reported by the investigator with the same time frame.

The AstraZeneca representative will work with the investigator to compile all the necessary information and ensure that all the necessary information is provided to Bristol-Myers Squibb Global Pharmacovigilance and Epidemiology within one calendar day of initial receipt for fatal and life threatening events and within three calendar days of initial receipt for all other SAEs.

All SAEs have to be reported, whether or not considered causally related to the investigational product, or to the study procedure(s). All SAEs will be recorded in the eCRF. SAEs will be recorded from the time of informed consent.

The investigator and/or Sponsor are responsible for informing the Ethics Committee (EC) about the SAE as per local requirements. Reporting of SAEs to the Regulatory Authority is the responsibility of Bristol-Myers Squibb.

An automated email alert will be sent to the designated AstraZeneca representative, when the page with SAE information is saved in WBDC system by the Investigators or other site personnel. If the WBDC system is not available, then the Investigator or other study site personnel reports by fax an SAE to the appropriate AstraZeneca representative. A paper back-

up SAE report is used for this purpose. The same reporting time frames still apply. The investigator is responsible for completing the eCRF as soon as the system becomes available again. The AstraZeneca representative will forward all information relevant to the SAE to Bristol-Myers Squibb Pharmacovigilance via fax or email.

#### **6.4.5 Laboratory safety assessment**

Blood and urine samples for determination of clinical chemistry, haematology and urinalysis will be taken at the times indicated in the Study Plan (See [Table 4](#) or [Table 5](#)).

The following laboratory variables will be measured. For blood volume see Section [7.1](#).

**Table 9 Safety laboratory variables**

Visit	1	2 <sup>a</sup>	3 <sup>a</sup>	4	5	6	7 <sup>b</sup>	8	9	10	11	12	13	14	15	16	17	18
Study week	-12 or -6	-10	-7	-4	-1	0	1	4	8	12	16	20	24	32	40	48	52	55
<b>Haematology</b>																		
Haemoglobin	X			X		X	X	X	X	X	X	X	X	X	X	X	X	X
Haematocrit	X			X		X	X	X	X	X	X	X	X	X	X	X	X	X
Red blood cell count	X			X		X	X	X	X	X	X	X	X	X	X	X	X	X
White blood cell count and differential	X			X		X	X	X	X	X	X	X	X	X	X	X	X	X
Platelet count	X			X		X	X	X	X	X	X	X	X	X	X	X	X	X
<b>Clinical chemistry (serum)</b>																		
AST (SGOT)	X			X		X	X	X	X	X	X	X	X	X	X	X	X	X
ALT (SGPT)	X			X		X	X	X	X	X	X	X	X	X	X	X	X	X
Alkaline Phosphatase (AP)	X			X		X	X	X	X	X	X	X	X	X	X	X	X	X
CK	X			X		X	X	X	X	X	X	X	X	X	X	X	X	X
Total Bilirubin (TB)	X			X		X	X	X	X	X	X	X	X	X	X	X	X	X
Blood Urea Nitrogen (BUN)	X			X		X	X	X	X	X	X	X	X	X	X	X	X	X
Electrolytes <sup>c</sup>	X			X		X	X	X	X	X	X	X	X	X	X	X	X	X
Total protein	X			X		X	X	X	X	X	X	X	X	X	X	X	X	X
Albumin	X			X		X	X	X	X	X	X	X	X	X	X	X	X	X
Uric acid	X			X		X	X	X	X	X	X	X	X	X	X	X	X	X
Serum Creatinine (SCr)	X			X		X	X	X	X	X	X	X	X	X	X	X	X	X
eGFR (Japanese guideline formula) <sup>d</sup>	X			X		X	X	X	X	X	X	X	X	X	X	X	X	X
Serum Cystatin C						X							X				X	
PTH						X							X				X	X
TSH	X																	

**Table 9 Safety laboratory variables**

Visit	1	2 <sup>a</sup>	3 <sup>a</sup>	4	5	6	7 <sup>b</sup>	8	9	10	11	12	13	14	15	16	17	18	
Study week	-12 or -6	-10	-7	-4	-1	0	1	4	8	12	16	20	24	32	40	48	52	55	
PT	X																		
PTT	X																		
Hepatitis Screen Panel <sup>c</sup>	X																		
<b>Urinalysis (dipstick)</b>																			
Glucose	X			X		X	X	X	X	X	X	X	X	X	X	X	X	X	
Blood <sup>f</sup>	X			X		X	X	X	X	X	X	X	X	X	X	X	X	X	
Pregnancy test <sup>g</sup>	X			X		X		X	X	X	X	X	X	X	X	X	X	X	
<b>Urinalysis (spot urine)</b>																			
Creatinine	X					X				X	X	X	X	X	X	X	X	X	
Albumin	X					X				X	X	X	X	X	X	X	X	X	
UACR	X					X				X	X	X	X	X	X	X	X	X	
Na	X					X				X	X	X	X	X	X	X	X	X	
Glucose	X					X				X	X	X	X	X	X	X	X	X	

a Visit 2 and 3 are only applicable for subgroup of monotherapy.

b Visit 7 (Week 1) is only applicable for subgroup of GLP-1 in combination therapy.

c Electrolytes: Sodium, Bicarbonate, Potassium, Chloride, Calcium, Magnesium, Phosphorus

d The eGFR will be estimated by the abbreviated Japanese guideline formula and reported by central laboratory.

e Includes anti-HAV (IgM, IgG), HBsAg, anti-HBc (IgG, IgM) and anti-HCV.

f Microscopy if dipstick positive for blood

g Urine HCG pregnancy test for WOCBP (HCG minimum sensitivity of 25 IU/L) (dipstick analyzed at the study centre)

#### **6.4.6 Physical examination**

- A brief physical examination should include the cardiovascular system, lungs, abdomen, and extremities, and any organ system pertinent to the subject's signs, symptoms, or AEs. The subject should always be evaluated for the presence of oedema.
- A complete physical examination should include general appearance, head, eyes, ears, nose, throat, neck, cardiovascular system, lungs, abdomen, lymph nodes, extremities, neurological system, skin, and musculoskeletal system. The subject should always be evaluated for the presence of oedema.

Baseline data is collected at Visit 6, and new findings discovered on subsequent physical examinations should be recorded as changes from baseline.

#### **6.4.7 ECG**

##### **Resting 12-lead ECG**

A 12-lead ECG will be taken after the subject has been lying down resting. The ECG will be evaluated by the investigator and entered as 'Normal' or 'Abnormal' in the eCRF. If the ECG is evaluated as "Abnormal" the investigator should document the specific abnormality.

#### **6.4.8 Vital signs**

##### **6.4.8.1 Pulse and blood pressure**

One pulse measurement will be taken after the subject has been sitting and resting for at least 5 minutes and before blood samples are taken. The pulse measurement will be followed by three BP measurements separated by 2 minutes each. All three BP readings should be recorded. At Visit 1 the seated BP will be recorded three times in both the left and the right arms. All three measurements should be made in one arm before transferring the cuff to the other arm. The arm with the highest mean seated BP readings will be the one used for all subsequent readings throughout the study. BP readings should be taken while the subject is in a comfortable seated position with the arm supported at the level of the heart. All readings should be recorded. Ideally, BP should be measured with the same machine, at the same time of day, and by the same personnel at each visit. A standard mercury sphygmomanometer with a standardised cuff adapted to the size of the subject's arm is recommended. Oscillometric devices (such as Dynamap) may be used at sites where:

- a mercury sphygmomanometer is not available, or
- a mercury sphygmomanometer is available, but site staff is not practiced in its use
- use of mercury devices is restricted by local law.



#### **6.4.8.2 Orthostatic blood pressure**

At selected visits where orthostatic BP and pulse are collected, supine and standing measurements should be made after the seated BP and pulse measurements have been made, using the same arm that was used for the seated BP measurements. All readings should be recorded. Ideally, BP should be measured with the same machine, at the same time of day, and by the same personnel at each visit.

##### **Supine BP and pulse**

The supine BP and pulse must be measured prior to the standing BP and pulse. After the subject rests in the supine position for at least 5 minutes, supine BP and pulse will be determined from three replicate measurements obtained at least 1 minute apart. All three readings must be recorded. For study analyses, the average of the three BP and pulse readings will be used.

##### **Standing BP and pulse**

After the supine BP and pulse measurements are obtained, the subject will stand for 2 to 3 minutes. After this time, the BP will be measured with the arm supported at the level of the heart. Standing BP and pulse will be determined from three replicate measurements obtained at least 1 minute apart. All 3 readings must be recorded. For study analyses, the average of the three BP and pulse readings will be used.

#### **6.4.9 Other safety assessments**

Self-monitored plasma glucose readings and hypoglycemic events will be collected in a subject diary and reviewed by the investigator at each visit.

##### **6.4.9.1 Fasting plasma glucose concentrations and hypoglycemic events**

Subject self-monitoring of FPG is performed in order to reduce the risks associated with prolonged hyperglycemia and to confirm symptoms of hypoglycemia. Subjects will be asked to perform self-monitoring of FPG using glucometers provided by AstraZeneca. The subjects will receive instructions for the use of the glucometer according to the manufacturer's instructions.

FPG should be self-monitored at least every second day between Visit 2 and 6 and at least once a week between Visit 6 and 8. The results should be recorded in the subject diary, which will be collected and reviewed by the study personnel at each visit starting with Visit 3; a print out will be stored in the investigator study file.

The memory of the glucometer should be reviewed and compared with the readings in the subject's diary. The glucose values should be reviewed by the study personnel to identify any unusually high or low values, and to confirm that self-monitoring was performed by the subject. If fingerstick glucose values are discordant with central laboratory results or with clinical symptoms, the subject's glucometer should be tested and the glucometer instructions should be re-reviewed with the subject.

If self-monitored FPG is above 270 mg/dL (15 mmol/L) from Week - 4 (Visit 4) to Week 4 (Visit 8), and above 240 mg/dL (13.2 mmol/L) from Week 4 (Visit 8) to Week 52 (Visit 17), the subject should repeat the FPG on the same day. If the second FPG value is above 270 mg/dL (15 mmol/L), or 240 mg/dL (13.2mmol/L), respectively, the subject should contact the study centre and be scheduled for a central laboratory FPG measurement within one week.

If central laboratory values show similar values, the subject should be discontinued.

A separate section in the eCRF will be used to document all reported episodes of hypoglycemia after Week 0 (Visit 6). The subjects will be asked to check their blood glucose if they develop symptoms suggestive of hypoglycemia and to record specific symptoms in the subject diary. The Investigator is responsible for questioning the subject about all symptoms reported in the diary and for determining if they meet the clinical definition of hypoglycemia. Only symptoms and/or blood glucose values that meet the definition of hypoglycemia should be reported on the hypoglycemia eCRF pages.

A hypoglycemic event can be either:

- Symptoms of hypoglycemia with a low blood glucose reading (<63 mg/dL [ $<3.5$  mmol/L])
- A low blood glucose reading (<63 mg/dL [ $<3.5$  mmol/L])
- Symptoms of hypoglycemia without a blood glucose reading

Hypoglycemic episodes or discontinuation due to hypoglycemia should not be reported on the AE eCRF page unless the event fulfils protocol criteria for a SAE (see Section 6.4.2). In this case, an SAE must be reported in addition to the hypoglycemia eCRF pages for hypoglycemia.

Symptoms suggestive of hypoglycemia with an associated capillary or plasma glucose value  $\geq 63$  mg/dL ( $\geq 3.5$  mmol/L), should be recorded as an AE rather than as a hypoglycemic event. If the physician does not consider the glucose measurement to be accurate, however, the event should be documented as a hypoglycemic event in the hypoglycemia eCRF.

For the evaluation of hypoglycemic events, this study will use the definitions provided in the CPMP guidance on clinical investigation of medicinal products in the treatment of diabetes mellitus, as described below.

- Major hypoglycemic events, defined as symptomatic events requiring external assistance due to severe impairment in consciousness or behaviour, with a capillary or plasma glucose value <54 mg/dL (<3.0 mmol/L), and prompt recovery after glucose or glucagon administration.
- Minor hypoglycemic event, defined as either a symptomatic episode with a capillary or plasma glucose measurement <63 mg/dL (<3.5 mmol/L) regardless of need for external assistance or an asymptomatic capillary or plasma glucose

measurement below 63 mg/dL (3.5 mmol/L) , that does not qualify as a major episode

- Events suggestive of hypoglycemia, defined as a symptomatic event without a confirmatory blood glucose measurement.

Data to be collected for each hypoglycemic event:

- Date and time of episode (start and stop)
- Whether the subject was sleeping at the time of the event
- Whether symptoms were present, and list of symptoms
- Possible contributing factors
- Whether a fingerstick value was obtained, and if so, the plasma glucose value
- Whether intervention was needed for recovery
- How the episode was treated
- Whether recovery was prompt after treatment
- Time of last anti-diabetic agents administration
- Time of last meal and its contents

The subject diary will be reviewed and the hypoglycemic event data will be transcribed into the eCRFs at each clinical visit. A new diary for the next period will be handed over to the subject if needed. If a major hypoglycemic event or more than one minor event has occurred since the previous visit, the subject should contact the investigator.

#### **6.4.9.2 Urinary and Genital Infections**

The following is presented to assist in the classification and management of infections of the urinary and genital tracts in studies with dapagliflozin. It is not intended to supplant investigators' clinical judgement.

##### **Urinary Tract Infections**

If based on the suggestive signs or symptoms (dysuria, urgency or frequency of urination, suprapubic or perineal discomfort, flank, back, or abdominal pain, costovertebral angle tenderness, nausea, vomiting, fever, chills, or sepsis) the investigator believes that a urinary tract infection may be present, urine cultures (in a local laboratory) should be obtained to confirm a presumptive diagnosis of cystitis, urinary tract infection, pyelonephritis, or prostatitis. Mid-stream clean catch urine collections are recommended. Clinical judgement and local standards of care should apply to decisions concerning therapy. It is strongly

recommended that a diagnosis of a recurrent urinary tract infection, defined either as two infections in a 6-month period or three infections in a 12-month period, should result in referral to a gynecologist (women) or urologist (men) for further diagnosis and treatment. Any treatment needs to be documented in the eCRF.

Between scheduled visits, subjects may experience novel signs or symptoms that are potentially indicative of urinary or a genital tract infection. The subject should contact the investigator by telephone. An unscheduled visit will be planned as soon as possible, preferably within 24 h. The investigator will take the subject's recent history, a midstream urine sample will be obtained for urine analyses and a mandated urine culture, and it is also recommended that a genital swab is done, if indicated. Analyses and culture(s) are to be performed at the local laboratory. The investigator will follow local guidelines to recommend treatment for urinary tract infection or genital tract infection.

Investigational product should be temporarily stopped in subjects with clinical evidence of upper urinary tract infection (eg, pyelonephritis) or presumed urosepsis until the course of treatment of the infection has been completed and clinical recovery has occurred. It is recommended that a follow-up urine culture is obtained within 7 days of clinical recovery from a documented UTI. Whether additional therapy is prescribed because of culture results should be determined by Investigator judgement.

### **Genital Tract Infections**

The diagnosis of vaginitis, vulvovaginitis, vulvitis or balanitis can be made based on physical examinations, culture of secretions or a therapeutic response to treatment of fungal or other vaginal pathogens. A urine culture is not required for diagnosis of genital infections if the diagnosis is confirmed by physical examination, culture of secretions, or a therapeutic response to treatment of fungal or other vaginal pathogens. A recurrent genital tract infection, defined as more than 2 infections in a 6 month period, should result in a referral to a gynecologist (women) or urologist (men) for further diagnosis and treatment. Any treatment needs to be documented in the eCRF.

Also, it is the investigator's responsibility to report, as applicable based on investigator's judgement and subject's medical history, related AEs as defined in Section 6.4.5. Additional information, including but not limited to completion of supplemental eCRFs may be requested for certain AEs and/or laboratory abnormalities which are reported/identified during the course of the study.

### **Asymptomatic bacteriuria**

During enrolment, treatment and follow up of subjects in this trial, the investigator may discover a subject with asymptomatic bacteriuria. Asymptomatic bacteriuria is defined as the presence of  $\geq 10^5$  colony forming units/mL of bacteria, in a properly collected voided urine specimen, without signs or symptoms typically attributed to urinary tract infection. Asymptomatic bacteriuria is prevalent among diabetic women, and is associated with pyuria in 70% of cases. Neither guidelines from the US (Nicolle et al 2005, USPST 2004) nor Europe (European Association of Urology 2008) recommend screening for, or treatment of,

asymptomatic bacteriuria in non-pregnant diabetic subjects. In this study, the central laboratory will report urinary dipstick test results for hemoglobin but will not routinely report the results of urinary dipstick tests for leukocyte esterase as a screening test for pyuria in surveillance urine examinations.

#### **6.4.10 Congestive Heart Failure**

The risk of electrolyte abnormalities, volume depletion, and impaired renal function is enhanced when two diuretics are used in combination. For this reason, caution should be exercised when administering dapagliflozin, which has a modest diuretic effect, to subjects who are taking loop diuretics. These subjects should have careful monitoring of electrolytes, volume status, and renal function. Loop diuretic dose adjustments should be made if clinically indicated.

#### **6.4.11 Change in kidney function**

Please see Appendix [F](#) for further guidance.

#### **6.4.12 Hyponatremia**

Please see Appendix [E](#) for further guidance.

#### **6.4.13 CK abnormalities**

Please see Section [5.8](#).

#### **6.4.14 Liver function test abnormalities**

Please see Section [5.8](#) and Appendix [G](#).

#### **6.4.15 Independent Adjudication Committee**

A Clinical Event Committee (CEC), blinded to the treatment of the subjects, will independently adjudicate certain cardiovascular AEs, and they will operate in accordance with a dedicated CEC Charter/Manual of Operations: Dapagliflozin Program. The CEC will adjudicate events possibly related to the following:

Death including:

1. Cardiovascular Death
2. Non-cardiovascular Death

Myocardial Infarction (MI) including:

1. ECG and /or cardiac enzymes confirmed MI
2. Sudden death
3. Percutaneous coronary intervention-related MI

4. Coronary artery bypass graft-related MI
5. MI diagnosed via pathologic criteria
6. Silent MI

Fatal and Non-fatal Stroke including:

1. Ischaemic Stroke
2. Haemorrhagic Stroke

SAEs of the following:

1. Heart failure
2. Cardiac arrhythmia
3. Unstable angina
4. Unplanned arterial revascularization (coronary, carotid and peripheral)
5. Cardiac arrest with successful resuscitation
6. Deep Vein Thrombosis and Pulmonary Emboli
7. Systemic non-stroke arterial embolism/thrombosis including systemic arterial occlusion
8. Non-traumatic amputation of the lower limb. Only events above the ankle will be considered for adjudication.

In order to provide the independent CEC with appropriate and adequate information for adjudication of the listed events, please consult the Reference Manual, Dapagliflozin Cardiovascular Adjudication Reference Manual for Primary Investigators and Study Staff.

#### **6.4.16 Hepatic Adjudication Committee**

An independent Hepatic Adjudication Committee will determine the probability that drug-induced liver injury (DILI) is the cause of liver-related abnormalities, including, but not limited to:

- Hepatic disorders leading to discontinuation from study treatment and/or death
- Liver laboratory abnormalities such as elevated AST and/or ALT with or without TB elevations.

A separate Adjudication Manual will define and describe the procedure for the handling, reporting and classification of these cases.

## **6.5 Patient reported outcomes (PRO) – Not applicable**

## **6.6 Pharmacokinetics – Not applicable**

## **6.7 Pharmacodynamics – Not applicable**

## **6.8 Pharmacogenetics – Not applicable**

## **6.9 Health economics – Not applicable**

# **7. BIOLOGICAL SAMPLING PROCEDURES**

## **7.1 Volume of blood**

The total volume of blood that will be drawn from each subject in this study is as follows:

**Table 10** Volume of blood to be drawn from each subject (mL)

Visit	1	2 <sup>a</sup>	3 <sup>a</sup>	4	5	6	7 <sup>b</sup>	8	9	10	11	12	13	14	15	16	17	18	Maximum volume
Sample volume	23	2	2	13.5	5	18.5	13.5	13.5	13.5	18.5	13.5	13.5	18.5	13.5	18.5	13.5	18.5	13.5	246

a Visit 2 and 3 are only applicable for subgroup of monotherapy.

b Visit 7 (Week 1) is only applicable for subgroup of GLP1 in combination therapy.

## **7.2 Handling, storage and destruction of biological samples**

The samples will be used up or disposed of after analyses.

## **7.3 Labelling and shipment of biohazard samples**

The Principal Investigator ensures that samples are labelled and shipped in accordance with the Laboratory Manual and the Biological Substance, Category B Regulations (materials containing or suspected to contain infectious substances that do not meet Category A criteria), see Appendix C ‘IATA 6.2 Guidance Document’.

Any samples identified as Infectious Category A materials are not shipped and no further samples will be taken from the subject unless agreed with AstraZeneca and appropriate labelling, shipment and containment provisions are approved.

## **7.4 Chain of custody of biological samples**

A full chain of custody is maintained for all samples throughout their lifecycle.

The Principal Investigator keeps full traceability of collected biological samples from the subjects while in storage at the centre until shipment or disposal (where appropriate) and keeps documentation of receipt of arrival.

The sample receiver keeps full traceability of the samples while in storage and during use until used or disposed of or until further shipment and keeps documentation of receipt of arrival.

AstraZeneca keeps oversight of the entire life cycle through internal procedures, monitoring of study sites and auditing of external laboratory providers.

Samples retained for further use are registered in the AstraZeneca biobank system during the entire life cycle.

## **7.5 Withdrawal of informed consent for donated biological samples**

If a subject withdraws consent to the use of donated biological samples, the samples will be disposed of/destroyed, and the action documented. If samples are already analysed, AstraZeneca is not obliged to destroy the results of this research.

The Principal Investigator:

- Ensures subjects' withdrawal of informed consent to the use of donated samples is notified immediately to AstraZeneca
- Ensures that biological samples from that subject, if stored at the study site, are immediately identified, disposed of /destroyed, and the action documented
- Ensures the laboratory(ies) holding the samples is/are informed about the withdrawn consent immediately and that samples are disposed/destroyed, the action documented and the signed document returned to the study site
- Ensures that the subject and AstraZeneca are informed about the sample disposal.

AstraZeneca ensures the central laboratory(ies) holding the samples is/are informed about the withdrawn consent immediately and that samples are disposed of/destroyed and the action documented and returned to the study site.

## **8. ETHICAL AND REGULATORY REQUIREMENTS**

### **8.1 Ethical conduct of the study**

The study will be performed in accordance with ethical principles that have their origin in the Declaration of Helsinki and are consistent with ICH/Good Clinical Practice (GCP), applicable regulatory requirements and the AstraZeneca policy on Bioethics and Human Biological Samples.



The applicable regulatory requirements in Japan are ‘Good Clinical Practice for Trials on Drugs (MHLW Ordinance No. 28, 27 March 1997, partially revised by MHLW Ordinance and their related notifications.

## **8.2 Subject data protection**

The Master Informed Consent Form will explain that: Study data will be stored in a computer database, maintaining confidentiality in accordance with national data legislation. Subject data will be maintaining confidentiality in accordance with national data legislation. For data verification purposes, authorised representatives of AstraZeneca, a regulatory authority, an EC may require direct access to parts of the hospital or practice records relevant to the study, including subjects’ medical history. All data computer processed by AstraZeneca will be identified by study code and enrolment code (E-code).

## **8.3 Ethics and regulatory review**

An EC should approve the final study protocol, including the final version of the Informed Consent Form and any other written information and/or materials to be provided to the subjects. The investigator/The Head of the study site will ensure the distribution of these documents to the applicable EC, and to the study site staff.

The opinion of the EC should be given in writing. The head of the study site should submit a notification of direction/determination as well as a copy of the institutional review board (IRB) written approval to AstraZeneca before enrolment of any subject should into the study.

The EC should approve all advertising used to recruit subjects for the study.

AstraZeneca should approve any modifications to the Informed Consent Form that are needed to meet local requirements.

The protocol should be re-approved by the IRB annually. The Principal Investigator should submit progress reports to the IRB via the head of the study site at the time of the protocol re-approval.

Before enrolment of any subject into the study, the final study protocol, including the final version of the Informed Consent Form, is approved by the national regulatory authority or a notification to the national regulatory authority is done, according to local regulations.

AstraZeneca will handle the distribution of any of these documents to the national regulatory authorities.

AstraZeneca will provide Regulatory Authorities, ECs, the head of the study site and Principal Investigators with safety updates/reports according to local requirements, including SUSARs (Suspected Unexpected Serious Adverse Reactions), where relevant.

The Head of the study site should submit a written report to the IRB providing the details of all AE case(s) reported by AstraZeneca.

## **8.4 Informed consent**

The Principal Investigator(s) at each centre will:

- Ensure each subject is given full and adequate oral and written information about the nature, purpose, possible risk and benefit of the study
- Ensure each subject is notified that they are free to discontinue from the study at any time
- Ensure that each subject is given the opportunity to ask questions and allowed time to consider the information provided
- Ensure each subject provides signed and dated informed consent before conducting any procedure specifically for the study
- Ensure the original, signed Informed Consent Form(s) is/are stored in the Investigator's Study File
- Ensure a copy of the signed Informed Consent Form is given to the subject
- Ensure that any incentives for subjects who participate in the study as well as any provisions for subjects harmed as a consequence of study participation are described in the informed consent form that is approved by an EC.

If any new information on the study medication becomes available which may influence the decision of the subject to continue the study, the investigator(s) should inform the subject of such information immediately, record this in a written form, and confirm with the subject if he or she wishes to continue the participation in the study. In addition, if the investigator(s) deem it necessary to revise the Informed Consent Form, they should revise it immediately (See Section 8.5). The investigator(s) should re-explain the subjects using updated Informed Consent Form even if although the subjects have already been informed of the new information verbally. Written informed consent to continue participation in the study should be provided separately.

## **8.5 Changes to the protocol and informed consent form**

Study procedures will not be changed without the mutual agreement of the Principal Investigator and AstraZeneca. If it is necessary for the study protocol to be amended, the amendment should be submitted to the Head of the Study Site and be approved by its IRB. If applicable, AstraZeneca should submit a notification to the regulatory authority before it is implemented. If a protocol amendment requires a change to a particular centre's Informed Consent Form, then AstraZeneca and the centre's IRB should be notified. Approval of the revised Informed Consent Form by AstraZeneca and by the IRB is required before the revised form is used. If an administrative change is required, such a change should be notified to or approved by each IRB according to local requirements.

## **8.6 Audits and inspections**

Authorised representatives of AstraZeneca, a regulatory authority, or an EC may perform audits or inspections at the centre, including source data verification. The purpose of an audit or inspection is to systematically and independently examine all study-related activities and documents, to determine whether these activities were conducted, and data were recorded, analysed, and accurately reported according to the protocol, GCP, guidelines of the International Conference on Harmonisation (ICH), and any applicable regulatory requirements. The investigator will contact AstraZeneca immediately if contacted by a regulatory agency about an inspection at the centre.

All study data may undergo a reliability review and onsite-GCP inspection by the regulatory authorities.

## **9. STUDY MANAGEMENT BY ASTRAZENECA**

### **9.1 Pre-study activities**

Before the first subject is entered into the study, it is necessary for a representative of AstraZeneca to visit the investigational study site to:

- Determine the adequacy of the facilities
- Determine availability of appropriate subjects for the study
- Discuss with the investigator(s) (and other personnel involved with the study) their responsibilities with regard to protocol adherence, and the responsibilities of AstraZeneca or its representatives.

### **9.2 Training of study site personnel**

Before the first subject is entered into the study, an AstraZeneca representative will review and discuss the requirements of the Clinical Study Protocol and related documents with the investigational staff and also train them in any study specific procedures and the WBDC system(s) utilised.

The Principal Investigator will ensure that appropriate training relevant to the study is given to all of these staff, and that any new information relevant to the performance of this study is forwarded to the staff involved.

The Principal Investigator will maintain a record of all individuals involved in the study (medical, nursing and other staff).

### **9.3 Monitoring of the study**

During the study, an AstraZeneca representative will have regular contacts with the study site, including visits to:

- Provide information and support to the investigator(s)
- Confirm that facilities remain acceptable
- Confirm that the investigational team is adhering to the protocol, that data are being accurately and timely recorded in the eCRFs, that biological samples are handled in accordance with the Laboratory Manual and that study drug accountability checks are being performed
- Perform source data verification (a comparison of the data in the eCRFs with the subject's medical records at the hospital or practice, and other records relevant to the study) including verification of informed consent of participating subjects. This will require direct access to all original records for each subject (eg, clinic charts)
- Ensure withdrawal of informed consent to the use of the subject's biological samples is reported and biological samples are identified and disposed of/destroyed accordingly, and the action is documented, and reported to the subject.

The AstraZeneca representative will be available between visits if the investigator(s) or other staff at the centre needs information and advice about the study conduct.

### **9.3.1 Source data**

Source data are any data generated as a result of the subject's inclusion in the study (including run-in and/or follow up related to the study) and includes all related medical examinations and other records.

Original data recorded on the eCRFs and regarded as source data are as follows.

- Reason for therapy or medication
- Any Comments on eCRF
- Evaluation for inclusion/exclusion criteria
- AE (Yes/No, Maximum intensity, Serious, Treatment required, Outcome, Causality)
- Description of AE on Serious AE form
- Main reason for premature discontinuation

### **9.3.2 Direct access to source data in Japan**

The Head of the institution and the Principal Investigator/sub-investigator will cooperate for monitoring and audit by AstraZeneca, and accept inspection by the IRB or regulatory authorities. All study documents such as raw data will be open for direct access to source data

at the request of the monitor and the auditor of AstraZeneca, the IRB, or regulatory authorities.

The monitor(s) will verify data from the eCRFs against source data before collecting the eCRFs to ensure accuracy and completeness of documentation, and assure that the Principal Investigator/sub-investigator has submitted the eCRFs to AstraZeneca. If the investigator wishes to amend the collected eCRFs, the monitor will ensure that the Principal Investigator/sub-investigator has documented the amendment in writing (signed and dated) and provided this to AstraZeneca.

## 9.4 Study agreements

The Principal Investigator at each/the centre should comply with all the terms, conditions, and obligations of the Clinical Study Agreement, or equivalent, for this study. In the event of any inconsistency between this Clinical Study Protocol and the Clinical Study Agreement, the terms of Clinical Study Protocol shall prevail with respect to the conduct of the study and the treatment of subjects and in all other respects, not relating to study conduct or treatment of subjects, the terms of the Clinical Study Agreement shall prevail.

Agreements between AstraZeneca and the Principal Investigator should be in place before any study-related procedures can take place, or subjects are enrolled.

### 9.4.1 Archiving of study documents

- (i) **Study files.** AstraZeneca will provide the Principal Investigator with a file in which to organise and retain all study-related documents. All study documents (including letters from AstraZeneca) should be retained in this file by the Principal Investigator. The monitor will regularly check the file to ensure that all relevant documents are retained. The contents of the file may be audited/inspected by AstraZeneca's auditor, regulatory authorities, or IRB.
- (ii) **Period of record retention.** The study site (and the Principal Investigator) will retain the essential documents specified in the ICH GCP (eg, source document such as medical records, contract, signed consent form). Essential documents should be retained at the study site for at least 15 years following completion of the study, or per regulatory obligations if longer, and thereafter destroyed only after agreement with AstraZeneca. However this is not always applied to those that are not preservable such as blood samples. In the event of any inconsistency between the above-mentioned contents and the contract with the study site, the contract shall prevail. These documents should be retained for a longer period however if needed by AstraZeneca, and the specific period and method of retention will be separately discussed between the study site and AstraZeneca. AstraZeneca should notify the head of the study site in writing when the study related records are no longer needed. The records should be managed by a responsible person appointed by the head of the study site.

#### **9.4.2 Deviation from the clinical study protocol in Japan**

The investigator(s) must not deviate from or make any changes to the protocol without documented agreement between the principal investigator and AstraZeneca K.K. or the IRB approval based on its deliberations. However, this shall not apply to cases where the deviation or change is necessary to avoid an immediate hazard to the subjects or for other compelling medical reasons, or where the changes involve only logistical or administrative aspects of the clinical trial (eg, changes to the organisation/structure of the sponsor, the name/department name of the medical institution, the address or phone number of the medical institution or the sponsor, the job title of the investigator, and monitors).

The investigator(s) should document any deviation from the protocol regardless of their reasons. Only when the protocol was not followed in order to avoid an immediate hazard to the subjects or for other medically compelling reason, the investigator should prepare and submit the records explaining the reasons thereof to the sponsor, and retain a copy of the records.

The investigator(s) may deviate from or make a change to the protocol without documented agreement between the principal investigator and AstraZeneca K.K. or the IRB approval, only in the event of a medical emergency, eg, it is only way to avoid an immediate hazard to the subjects. In such case, the principal investigator must notify details of the deviation or change, the reason, and a proposed revision in the protocol if required, to AstraZeneca K.K. and the head of the study site and IRB via the head of the study site as soon as possible, in order to obtain their approval. A certificate of approval by the head of the study site as well as AstraZeneca K.K. should be obtained via the head of the study site.

#### **9.5 Study timetable and end of study**

The end of the study is defined as ‘the last visit of the last subject undergoing the study’.

The study may be terminated at individual centres if the study procedures are not being performed according to GCP, or if recruitment is slow. AstraZeneca may also terminate the entire study prematurely if concerns for safety arise within this study or in any other study with dapagliflozin.

Planned duration of the study:

Study period: February 2011 - October 2012

Registration period: February 2011 - October 2011

#### **Discontinuation or suspension of the whole study programme**

If AstraZeneca decides to prematurely terminate or suspend the study, the Principal Investigator, the head of the institution, and regulatory authorities should receive written notification of the reasons for the premature termination or suspension.

The Principal Investigator will immediately notify the decision to the subjects, give appropriate medical treatment; take necessary measures, and record treatment or measures provided on the source documents.

### **Completion of the study**

Upon terminating the study, the Principal Investigator will report in writing the completion of the study as well as the summary of the results to the head of the study site in accordance with the institution's rules. The head of the study site, who is informed of the termination by the investigator, will provide a written notification of the results to the IRB and AstraZeneca.

## **10. DATA MANAGEMENT BY ASTRAZENECA DATA MANAGEMENT CENTRE**

Data management will be performed by AstraZeneca Data Management Centre staff.

Data will be entered in the WBDC system at the study site. Trained study personnel will be responsible for entering data on the observations, tests and assessments specified in the protocol into the WBDC system and according to the eCRF Instructions. The eCRF Instructions will also guide the study site in performing data entry.

Data entered in the WBDC system will be immediately saved to a central database and changes tracked to provide an audit trail. The data will then be Source Data Verified (SDV), reviewed, queried and updated as needed. Principal Investigator is responsible for signing the eCRF electronically as per the eCRF instructions.

Data queries will be raised for inconsistent, impossible or missing data. All entries to the study database will be available in an audit trail. Quality assurance procedures will be applied to each stage of data handling to ensure that all data are reliable and have been processed correctly.

The data will be frozen and then locked to prevent further editing. When all data have been coded, validated, signed and locked, clean file will be declared. All editing accesses will be revoked and the final database will be locked. Copy of the eCRF will be archived at the study site when the study has been locked.

The study Data Management Plan will describe in greater detail the methods used to collect, check and process clinical data. It will also clarify the roles and responsibilities of the various functions and personnel involved in the data management process.

Prior to database lock, all decisions on the evaluability of the data from each individual subject must have been made and documented.

## **Management of external data**

Data Management determines the format of the data to be received from external vendors and coordinates the flow of data to an external environment or clinical database (if applicable). Data Management will ensure that the data collection tool (eg, IWRS etc) will be tested / validated as needed. External data reconciliation will be done with the clinical database as applicable.

## **Dictionary coding**

AEs and medical/surgical history will be classified according to the terminology of the latest version the MedDRA. Medications will be classified according to the Bristol-Myers Squibb Drug Dictionary. All coding will be performed by the coding team at Bristol-Myers Squibb.

# **11. EVALUATION AND CALCULATION OF VARIABLES BY ASTRAZENECA**

## **11.1 Calculation or derivation of efficacy variable(s)**

### **11.1.1 Change and percent change from baseline**

Change from baseline will be calculated as absolute difference (or percentage change) between the value measured at or derived for a specific time point after baseline minus baseline value. Baseline is defined as the day of first administration of dapagliflozin, and the baseline value is the last assessment on the day of first administration of dapagliflozin but also before during intake.

Percent change from baseline will be computed as  $100 * (\text{value measured at or derived for a specific time point after baseline} - \text{baseline value}) / \text{baseline value}$ .

### **11.1.2 Last observation carried forward (LOCF)**

If no measurement is available at a time point, the last post-baseline measurement prior to the specific time-point will be used instead for analysis. Unless otherwise specified, if a subject initiates rescue medication, the last value taken on or before the first rescue will be used for analysis.

## **11.2 Calculation or derivation of safety variable(s)**

### **11.2.1 Calculation of eGFR**

The following Japan guideline recommended equation will be used to calculate eGFR.

$$\text{eGFR (mL/min/1.73m}^2\text{)} = 194 \times (\text{SCr})^{-1.094} \times (\text{age})^{-0.287} \times (0.739 \text{ if female})$$

### **11.2.2 Other significant adverse events (OAE)**

During the evaluation of the AE data, an AstraZeneca medically qualified expert will review the list of AEs that were not reported as SAEs and Discontinuation of Investigational Product



due to Adverse Events (DAEs). Based on the expert's judgement, significant AEs of particular clinical importance may, after consultation with the Global Patient Subject Safety Physician, be considered OAEs and reported as such in the clinical study report. A similar review of laboratory/vital signs/ECG data will be performed for identification of OAEs.

Examples of these are marked haematological and other laboratory abnormalities, and certain events that lead to intervention (other than those already classified as serious), dose reduction or significant additional treatment.

- 11.3 Calculation or derivation of patient reported outcome variables – Not applicable**
- 11.4 Calculation or derivation of pharmacokinetic variables – Not applicable**
- 11.5 Calculation or derivation of pharmacodynamic variable(s) – Not applicable**
- 11.6 Calculation or derivation of pharmacogenetic variables – Not applicable**
- 11.7 Calculation or derivation of health economic variables – Not applicable**

## **12. STATISTICAL METHODS AND SAMPLE SIZE DETERMINATION BY ASTRAZENECA**

A comprehensive statistical analysis plan will be finalized prior to the clean file.

### **12.1 Description of analysis sets**

Analysis of safety variables will be based on safety analysis set. Full analysis set will be used for the analysis of efficacy variables.

#### **12.1.1 Full analysis set**

The full analysis set will include all subjects who received at least one dose of study medication, and who have a non-missing baseline value and at least one post-baseline efficacy value for at least one efficacy variable.

#### **12.1.2 Safety analysis set**

The safety analysis set will include all subjects who received at least one dose of study medication and who provide any safety records.

## **12.2 Methods of statistical analyses**

### **12.2.1 Demographics and baseline characteristics**

Descriptive summaries of demographics and baseline characteristics will be presented by subgroup and for all subjects combined using the safety analysis set.

### **12.2.2 Efficacy analysis**

Unless otherwise specified, for all changes (or percent changes) from baseline to a specific time point post-baseline as well as for glycemic therapeutic response definitions, analyses will be based on measurements available at that time point or the last post baseline measurement prior to the time-point, if no measurement is available at that time point, ie, LOCF. For rescued subjects, values collected after initiation of rescue medication will not be considered in calculating the LOCF values.

Descriptive statistics at each time point, including Week 52 (LOCF) will be presented by subgroup as well as for total population for all efficacy variables including and excluding data after rescue. Descriptive summaries will similarly be made by final dose.

### **12.2.3 Safety analysis**

The safety evaluations will include analyses of AEs, laboratory values, ECG, vital signs (pulse and BP), hypoglycemic events, eGFR and physical examination findings.

The analysis of safety will be based on the safety analysis set. Safety data gained during the 52-week treatment period as well as during the 3-week safety follow-up period will be evaluated. Safety data will be presented by subgroup as well as for total population. Safety data will also be presented by final dose. Safety variables will be summarized descriptively.

## **12.3 Determination of sample size**

Total 700 Japanese subjects with T2DM who have inadequate glycemic control on diet and exercise or on concomitant anti-diabetic agents is planned, with number of subjects in each subgroup as follows:

- Monotherapy: 240 subjects.
- Combination therapies with SU: 120 subjects.
- Combination therapies with DPP-4, AGI, metformin, or TZD: 60 subjects for each.
- Combination therapy with glinides or GLP-1: 50 subjects for each.

The number of subjects were determined in consideration of [Japanese Guideline on Clinical Evaluation methods on Oral Hypoglycemic Agents](#) suggesting at least 100 subjects each, for monotherapy and combination therapies considered at higher risk of hypoglycemia and at least 50 subjects each, for other combination therapies. 10% marginal subjects are added considering drop-out through the study.

## 12.4 Data monitoring committee – Not applicable

## 13. IMPORTANT MEDICAL PROCEDURES TO BE FOLLOWED BY THE INVESTIGATOR

### 13.1 Medical emergencies and AstraZeneca contacts

The Principal Investigator is responsible for ensuring that procedures and expertise are available to handle medical emergencies during the study. **A medical emergency usually constitutes an SAE and is to be reported as such, see Section 6.4.4**

In the case of a medical emergency the investigator may contact the Study Delivery Team Leader. If the Study Delivery Team Leader is not available, contact the Study Delivery Team Physician.

Name	Role in the study	Address & telephone number
	Study Delivery Team Leader	
	Study Delivery Team Physician	
Monitor	Study Delivery Team Monitor	See Supplement A, “Investigators and Study Administrative Structure”.

### 13.2 Overdose

Overdose is defined as the accidental or intentional ingestion of any dose of investigational product that is considered both excessive and medically important. Dapagliflozin has been well tolerated at doses of up to 500 mg/day in single dose testing in healthy volunteers and up to 100 mg/day in repeat dose testing for 14 days in healthy volunteers and subjects with T2DM. Once an investigator decides that a particular occurrence is an overdose, it must be reported as a SAE. If an overdose is suspected, monitoring of vital functions as well as treatment as appropriate should be performed.

### 13.3 Pregnancy

All outcomes of pregnancy must be reported to AstraZeneca.

Pregnancy itself is not regarded as an AE unless there is a suspicion that the investigational product under study and/or concomitant medication which defined in this protocol may have interfered with the effectiveness of a contraceptive medication. Congenital abnormalities/birth defects and spontaneous miscarriages should be reported and handled as SAEs. Elective abortions without complications should not be handled as AEs. The outcome of all pregnancies (spontaneous miscarriage, elective termination, normal birth or congenital abnormality) must be followed up and documented even if the subject was discontinued from the study.

If any pregnancy occurs in the course of the study, the subject should be discontinued, the investigational product should be stopped and then investigators or other site personnel must inform appropriate AstraZeneca representatives immediately but no later than the end of the next business day of when he or she becomes aware of it.

The PREGREP module in the eCRF is used to report the pregnancy. This module in the eCRF should be completed by the investigator and the AstraZeneca representative will forward the information to Bristol Myers Squibb using the same procedure as for SAE reporting. An AstraZeneca paper Pregnancy Outcome Report, part 2, is used to report the outcome of the pregnancy.

## **14. LIST OF REFERENCES**

### **ACCORD 2008**

The Action to Control Cardiovascular Risk in Diabetes Study Group. Effects of intensive glucose lowering in type 2 diabetes. *N Engl J Med* 2008; 358:2545-2559.

### **ADA 2009**

American Diabetes Association. Standards of medical care in diabetes- 2009. *Diabetes Care* 2009;32(supplement 1): S13-S61.

### **Bailey 2010**

Bailey CJ, Gross JL, Pieters A, Bastien A, List JF. Effect of dapagliflozin in patients with type 2 diabetes who have inadequate glycaemic control with metformin: a randomised, double-blind, placebo-controlled trial. *The Lancet* 2010;375:2223-33.

### **European Association of Urology 2008**

European Association of Urology. Guidelines on the management of urinary and male genital tract infections 2008. Available at <http://www.uroweb.org/>

### **Ferrannini, 2010**

Ferrannini E, Jimenez-Ramos S, Salsali A, Tang W, List JF. Dapagliflozin monotherapy in type 2 diabetes patients with inadequate glycaemic control by diet and exercise: a randomized, double-blind, placebo-controlled, phase III trial. *Diabetes Care* 2010, in press. (published online June 21, 2010)

**Gillett MJ, 2009**

Gillett MJ. International Expert Committee Report on the Role of the A1C Assay in the Diagnosis of Diabetes. *Clin Biochem Rev* 2009; 30: 197-200

**Japanese Guideline on Clinical Evaluation methods on Oral Hypoglycemic Agents**

**Koro et al 2004**

Koro CE, Bowlin SJ, Bourgeois N, Fedder DO. Glycaemic control from 1988 to 2000 among US adults diagnosed with type 2 diabetes: a preliminary report. *Diabetes Care* 2004; 27:17-20.

**Matthaei et al 2000**

Matthaei S, Stumvoll M, Kellner M, Haring HU. Pathophysiology and pharmacological treatment of insulin resistance. *Endocr Rev* 2000; 21:585-618.

**Meier and Butler 2005**

Meier JJ, Butler PC. Insulin secretion. In: DeGroot LJ, Jameson JL, editors. *Endocrinology*. 5 ed. Philadelphia: Elsevier Saunders; 2005. 961-73.

**Nathan 2006**

Nathan DM. Management of Hyperglycaemia in Type 2 Diabetes: A Consensus Algorithm for the Initiation and Adjustment of Therapy. *Diabetes Care* 2006; 29:1963-72.

**Nicolle et al 2005**

Nicolle LE, Bradley S, Colgan R, Rice JC, Schaeffer A, Hooton TM. Infectious diseases society of America guidelines for the diagnosis and treatment of asymptomatic bacteriuria in adults. *Clin Infect Dis* 2005; 40:643-54.

**Santer et al 2003**

Santer R et al. Molecular analysis of the SGLT2 gene in patients with renal glucosuria. *J Am Soc Nephrol* 2003; 14:2873-82.

**Scholl-Burgi et al 2004**

Scholl-Burgi et al. Long-term outcome of renal glucosuria type 0: the original patient and his natural history. *Nephrol Dial Transplant* 2004; 19:2394-6.

**Silverman 1991**

Silverman M. Structure and function of hexose transporters. *Annu Rev Biochem*. 1991; 60:757-94.

**UKPDS group 1998**

UK Prospective Diabetes Study (UKPDS) Group. Intensive blood-glucose control with sulphonylureas or insulin compared with conventional treatment and risk of complications in patients with type 2 diabetes (UKPDS 33). *Lancet*. 1998; 352:837-53.

Clinical Study Protocol  
Drug Substance Dapagliflozin  
Study Code D1692C00012  
Edition Number 1  
Date

**USPST 2004**

US Preventative Services Taskforce. Recommendation Statement Screening for Asymptomatic Bacteriuria. Revised 2004. Available at <http://www.ahrq.gov/clinic/uspstf/uspsbact.htm>

**Wright 2001**

Wright EM. Renal Na(+)-glucose cotransporters. *Am J Physiol Renal Physiol.* 2001; 280(1): F10-8.



---

**Clinical Study Protocol Appendix A**

Drug Substance      Dapagliflozin

Study Code          D1692C00012

Edition Number      1

Date

Protocol Dated

---

---

**Appendix A**  
**Signatures**

---

## ASTRAZENECA SIGNATURE(S)

---

**A long term open label study to evaluate the safety and efficacy of dapagliflozin as monotherapy or combination therapies with anti-diabetic drugs in Japanese patients with type 2 diabetes who have inadequate glycemic control**

---

This Clinical Study Protocol has been subjected to an internal AstraZeneca peer review.

I agree to the terms of this study protocol.

AstraZeneca Research and Development  
site representative



Shamik Parikh, MD  
Medical Science Director  
AstraZeneca Pharmaceuticals  
1800 Concord Pike  
Wilmington, DE  
19850-5437  
USA

12<sup>th</sup> Jan 2011  
Date  
(Day Month Year)

This document contains confidential information, which should not be copied, referred to, released or published without written approval from AstraZeneca. Investigators are cautioned that the information in this protocol may be subject to change and revision.



## ASTRAZENECA SIGNATURE(S)

---

**A long term open label study to evaluate the safety and efficacy of dapagliflozin as monotherapy or combination therapies with anti-diabetic drugs in Japanese patients with type 2 diabetes who have inadequate glycemic control**

---

This Clinical Study Protocol has been subjected to an internal AstraZeneca peer review.

I agree to the terms of this study protocol.

**AstraZeneca Research and Development**  
site representative

*Yasunari Taketani*  
Yasunari Taketani  
Study Delivery Team Leader  
AstraZeneca K.K.  
1-88, 1-chome, Ohyodo-naka, Kita-ku,  
Osaka 531-0076, Japan

*11 Jan, 2011*  
Date  
(Day Month Year)

This document contains confidential information, which should not be copied, referred to, released or published without written approval from AstraZeneca. Investigators are cautioned that the information in this protocol may be subject to change and revision.

## ASTRAZENECA SIGNATURE(S)

---


**A long term open label study to evaluate the safety and efficacy of dapagliflozin as monotherapy or combination therapies with anti-diabetic drugs in Japanese patients with type 2 diabetes who have inadequate glycemic control**

---

This Clinical Study Protocol has been subjected to an internal AstraZeneca peer review.

I agree to the terms of this study protocol.

AstraZeneca Research and  
Development site representative

  
\_\_\_\_\_  
Jennifer Sugg

Global Product Statistician  
AstraZeneca Pharmaceuticals LP  
Biostatistics  
USA

14 Jan 2011

Date  
(Day Month Year)

This document contains confidential information, which should not be copied, referred to, released or published without written approval from AstraZeneca. Investigators are cautioned that the information in this protocol may be subject to change and revision.

## ASTRAZENECA SIGNATURE(S)

---

**A long term open label study to evaluate the safety and efficacy of dapagliflozin as monotherapy or combination therapies with anti-diabetic drugs in Japanese patients with type 2 diabetes who have inadequate glycemic control**

---

This Clinical Study Protocol has been subjected to an internal AstraZeneca peer review.

I agree to the terms of this study protocol.

AstraZeneca Research and  
Development site representative

麻生 博也

Hiroya Asou  
Study Delivery Team Physician  
AstraZeneca K.K.

1-88, 1-chome, Ohyodo-naka, Kita-ku, Osaka  
531-0076, Japan

11th January, 2011

Date  
(Day Month Year)

This document contains confidential information, which should not be copied, referred to, released or published without written approval from AstraZeneca. Investigators are cautioned that the information in this protocol may be subject to change and revision.



---

**Clinical Study Protocol Appendix B**

Drug Substance	Dapagliflozin
Study Code	D1692C00012
Edition Number	1
Date	

---

---

**Appendix B**  
**Additional Safety Information**

---

## **FURTHER GUIDANCE ON THE DEFINITION OF A SERIOUS ADVERSE EVENT (SAE)**

### **Life threatening**

‘Life-threatening’ means that the subject was at immediate risk of death from the AE as it occurred or it is suspected that use or continued use of the product would result in the subject’s death. ‘Life-threatening’ does not mean that had an AE occurred in a more severe form it might have caused death (eg, hepatitis that resolved without hepatic failure).

### **Hospitalisation**

Outpatient treatment in an emergency room is not in itself a serious AE, although the reasons for it may be (eg, bronchospasm, laryngeal oedema). Hospital admissions and/or surgical operations planned before or during a study are not considered AEs if the illness or disease existed before the subject was enrolled in the study, provided that it did not deteriorate in an unexpected way during the study.

### **Important medical event or medical intervention**

Medical and scientific judgement should be exercised in deciding whether a case is serious in situations where important medical events may not be immediately life threatening or result in death, hospitalisation, disability or incapacity but may jeopardize the subject or may require medical intervention to prevent one or more outcomes listed in the definition of serious. These should usually be considered as serious.

Simply stopping the suspect drug does not mean that it is an important medical event; medical judgement must be used.

Examples of such events are:

- Angioedema not severe enough to require intubation but requiring iv hydrocortisone treatment
- Hepatotoxicity caused by paracetamol (acetaminophen) overdose requiring treatment with N-acetylcysteine
- Intensive treatment in an emergency room or at home for allergic bronchospasm
- Blood dyscrasias (eg, neutropenia or anaemia requiring blood transfusion, etc) or convulsions that do not result in hospitalisation
- Development of drug dependency or drug abuse

## **A GUIDE TO INTERPRETING THE CAUSALITY QUESTION**

The following factors should be considered when deciding if there is a “reasonable possibility” that an AE may have been caused by the drug.

- Time Course. Exposure to suspect drug. Has the subject actually received the suspect drug? Did the AE occur in a reasonable temporal relationship to the administration of the suspect drug?
- Consistency with known drug profile. Was the AE consistent with the previous knowledge of the suspect drug (pharmacology and toxicology) or drugs of the same pharmacological class? OR could the AE be anticipated from its pharmacological properties?
- Dechallenge experience. Did the AE resolve or improve on stopping or reducing the dose of the suspect drug?
- No alternative cause. The AE cannot be reasonably explained by another aetiology such as the underlying disease, other drugs, other host or environmental factors.
- Rechallenge experience. Did the AE reoccur if the suspected drug was reintroduced after having been stopped? AstraZeneca would not normally recommend or support a rechallenge.
- Laboratory tests. A specific laboratory investigation (if performed) has confirmed the relationship?

A “reasonable possibility” could be considered to exist for an AE where one or more of these factors exist.

In contrast, there would not be a “reasonable possibility” of causality if none of the above criteria apply or where there is evidence of exposure and a reasonable time course but any dechallenge (if performed) is negative or ambiguous or there is another more likely cause of the AE.

In difficult cases, other factors could be considered such as:

- Is this a recognised feature of overdose of the drug?
- Is there a known mechanism?

Ambiguous cases should be considered as being a “reasonable possibility” of a causal relationship unless further evidence becomes available to refute this. Causal relationship in cases where the disease under study has deteriorated due to lack of effect should be classified as no reasonable possibility.



---

**Clinical Study Protocol Appendix C**

Drug Substance	Dapagliflozin
Study Code	D1692C00012
Edition Number	1
Date	

---

---

**Appendix C  
International Airline Transportation Association (IATA) 6.2 Guidance  
Document**

---

## **LABELLING AND SHIPMENT OF BIOHAZARD SAMPLES**

International Airline Transportation Association (IATA) classifies biohazardous agents into 3 categories ([http://www.iata.org/whatwedo/cargo/dangerous\\_goods/infectious\\_substances.htm](http://www.iata.org/whatwedo/cargo/dangerous_goods/infectious_substances.htm)). For transport purposes the classification of infectious substances according to risk groups was removed from the Dangerous Goods Regulations (DGR) in the 46th edition (2005). Infectious substances are now classified either as Category A, Category B or Exempt. There is no direct relationship between Risk Groups and categories A and B.

**Category A Infectious Substances** are infectious substances in a form that, when exposure to it occurs, is capable of causing permanent disability, life-threatening or fatal disease in otherwise healthy humans or animals. Category A pathogens are eg, Ebola, Lassa fever virus:

- are to be packed and shipped in accordance with IATA Instruction 602.

**Category B Infectious Substances** are infectious Substances that do not meet the criteria for inclusion in Category A. Category B pathogens are eg, Hepatitis A, B, C, D, and E viruses, Human immunodeficiency virus (HIV) types 1 and 2. They are assigned the following UN number and proper shipping name:

- UN 3373 – Biological Substance, Category B
- are to be packed in accordance with UN3373 and IATA 650

**Exempt** - all other materials with minimal risk of containing pathogens

- Clinical trial samples will fall into Category B or exempt under IATA regulations
- Clinical trial samples will routinely be packed and transported at ambient temperature in IATA 650 compliant packaging ([http://www.iata.org/whatwedo/cargo/dangerous\\_goods/infectious\\_substances.htm](http://www.iata.org/whatwedo/cargo/dangerous_goods/infectious_substances.htm))
- **Biological samples transported in dry ice require additional dangerous goods specification for the dry-ice content**
- IATA compliant courier and packaging materials should be used for packing and transportation and packing should be done by an IATA certified person, as applicable



- Samples routinely transported by road or rail are subject to local regulations which require that they are also packed and transported in a safe and appropriate way to contain any risk of infection or contamination by using approved couriers and packaging / containment materials at all times. The IATA 650 biological sample containment standards are encouraged wherever possible when road or rail transport is used.



---

**Clinical Study Protocol Appendix D**

Drug Substance	Dapagliflozin
Study Code	D1692C00012
Edition Number	1
Date	

---

---

**Appendix D**  
**New York Heart Association (NYHA) Classification**

---

## **NEW YORK HEART ASSOCIATION (NYHA) CLASSIFICATION**

The NYHA classification will be based on the following definitions:

- Class I            No limitation:  
  
                         Ordinary physical exercise does not cause undue fatigue, dyspnoea or palpitations.
- Class II            Slight limitation of physical activity:  
  
                         Comfortable at rest but ordinary activity results in fatigue, palpitations, dyspnoea.
- Class III           Marked limitation of physical activity:  
  
                         Comfortable at rest but less than ordinary activity results in symptoms.
- Class IV            Unable to carry out any physical activity without discomfort:  
  
                         Symptoms of congestive heart failure are present even at rest with increased discomfort with any physical activity.



---

**Clinical Study Protocol Appendix E**

Drug Substance	Dapagliflozin
Study Code	D1692C00012
Edition Number	1
Date	

---

---

**Appendix E**  
**Algorithm on Management of Hyponatraemia**

---

## ALGORITHM ON MANAGEMENT OF HYPONATRAEMIA

If a patient experiences a serum sodium  $\leq 125$  mmol/L, with or without symptoms, dosing of investigational product will be interrupted. A repeat serum sodium concentration will be drawn within 3 days of the receipt of the result.

- If the repeat sodium concentration within 3 days is  $\geq 130$  mmol/L  
  
Investigational product may be restarted unless otherwise contraindicated. Serum sodium will be rechecked in 7 days after restarting the investigational product.
  - If the repeat sodium concentration within 7 days of restarting the investigational product is  $< 130$  mmol/L, investigational product will be discontinued. Patients will have an end of treatment visit and a follow-up visit three weeks after discontinuation of investigational product as specified in Section 5.8.2 in the CSP.
  - If the repeat sodium concentration within 7 days of restarting the investigational product is  $\geq 130$  mmol/L, further management should be based on composite of sodium concentration, clinical assessment of the patient and an evaluation of underlying cause of hyponatraemia.
- If the repeat sodium concentration within 3 days is  $< 130$  mmol/L  
  
If there is **no** suspected new, temporary, and reversible cause of hyponatraemia based on clinical assessment (other than investigational product), investigational product will be discontinued. Patients will have an end of treatment visit and a follow-up visit three weeks after discontinuation of investigational product as specified in Section 5.8.2 in the CSP.  
  
If there is a suspected new, temporary, and reversible cause of hyponatraemia based on clinical assessment (other than investigational product), investigational product will continue to be interrupted. The suspected cause of hyponatraemia should be identified and corrected. The serum sodium will be rechecked in another 7 days.
  - If the repeat sodium concentration within 7 days is  $< 130$  mmol/L, investigational product will be discontinued. Patients will have an end of treatment visit and a follow-up visit three weeks after discontinuation of investigational product as specified in Section 5.8.2 in the CSP.
  - If the repeat sodium level concentration within 7 days is  $\geq 130$  mmol/L investigational product may be restarted unless otherwise contraindicated. Serum sodium will be rechecked in 7 days after restarting the investigational product, and further management should be based on composite of sodium

concentration, clinical assessment of the patient and an evaluation of underlying cause of hyponatraemia.

For patients whose serum sodium is in the range of 126-129 mmol/L, the investigator's clinical judgment should apply concerning whether such patients should be entered into this algorithm.



---

**Clinical Study Protocol: Appendix F**

Drug Substance	Dapagliflozin
Study Code	D1692C00012
Edition Number	1
Date	

---

---

**Appendix F**  
**Case Identification and Management of Decreased Renal Function**

---

## **CASE IDENTIFICATION AND MANAGEMENT OF DECREASED RENAL FUNCTION**

In order to standardise the definition and management of decreased renal function, the following guideline has been developed. Please note separate sections for patients who are treated with metformin and for those who are not treated with metformin.

### FOR PATIENTS WHO ARE TREATED WITH METFORMIN:

**If calculated eGFR is <60 ml/min or if there is an increase in serum creatinine of  $\geq 0.5$  mg/dL ( $\geq 44.2$   $\mu\text{mol/L}$ ) above the baseline value based on central laboratory results, this will be considered a case of “decreased renal function”.**

In this circumstance, the investigator should consider evaluating the patient for potentially reversible causes of renal dysfunction including:

1. concurrent use of NSAIDS, antibiotics, or other medications known to affect measures of serum creatinine
2. volume depletion
3. urinary tract infection
4. obstructive uropathy.

Patients should return for repeat central laboratory testing as soon as possible, no later than 7 days after the abnormal result. Investigational drug and metformin should be interrupted pending the results of repeat testing. If after interruption eGFR is still <60 ml/min or if the serum creatinine value remains  $\geq 0.5$  mg/dL ( $\geq 44.2$   $\mu\text{mol/L}$ ) above the baseline value, the patient should permanently discontinue the study medication and be withdrawn from the study (in which case an Adverse Event must be reported).

If after interruption of investigational drug and metformin eGFR is  $\geq 60$  ml/min or if the serum creatinine value has decreased to <0.5 mg/dL (<44.2  $\mu\text{mol/L}$ ) above baseline, investigational drug and metformin can be re-started if appropriate in the judgement of the investigator and following consultation with the study team physician. If the patient restarts investigational drug and metformin and eGFR is again <60 ml/min or the serum creatinine value increases again to  $\geq 0.5$  mg/dL ( $\geq 44.2$   $\mu\text{mol/L}$ ) above baseline, the patient should permanently discontinue the study medication and be withdrawn from the study (in which case an Adverse Event must be reported).

### FOR PATIENTS WHO ARE NOT TREATED WITH METFORMIN:

**If a) eGFR is <45 mL/min or b) in subjects with baseline creatinine  $\geq 1.4$  mg/dL ( $\geq 123$   $\mu\text{mol/L}$ ) there is an absolute increase in serum creatinine of  $\geq 1.0$  mg/dL ( $\geq 88$   $\mu\text{mol/L}$ ), or c) in subjects with baseline creatinine <1.4 mg/dL (<123  $\mu\text{mol/L}$ ) there**



**is an absolute increase in serum creatinine of  $\geq 0.5$  mg/dL ( $\geq 44$   $\mu$ mol/L), this will be considered a case of “decreased renal function”.**

In this circumstance, the investigator should consider evaluating the patient for potentially reversible causes of renal dysfunction including:

1. concurrent use of NSAIDS, antibiotics, or other medications known to affect measures of serum creatinine
2. volume depletion
3. urinary tract infection
4. obstructive uropathy.

Patients should return for repeat central laboratory testing as soon as possible, no later than 7 days after the abnormal result. Investigational drug should be interrupted pending the results of repeat testing. If after interruption eGFR is still  $< 45$  ml/min or if the serum creatinine value remains  $\geq 0.5$  mg/dL ( $\geq 44.2$   $\mu$ mol/L) above the baseline value, the patient should permanently discontinue the study medication and be withdrawn from the study (in which case an Adverse Event must be reported).

If after interruption of investigational drug eGFR is  $\geq 45$  ml/min or if the serum creatinine value has decreased to  $< 0.5$  mg/dL ( $< 44.2$   $\mu$ mol/L) above baseline, investigational drug and metformin can be re-started if appropriate in the judgement of the investigator and following consultation with the study team physician. If the patient restarts investigational drug and eGFR is again  $< 45$  ml/min or the serum creatinine value increases again to  $\geq 0.5$  mg/dL ( $\geq 44.2$   $\mu$ mol/L) above baseline, the patient should permanently discontinue the study medication and be withdrawn from the study (in which case an Adverse Event must be reported).

---

**Clinical Study Protocol Appendix G**

Drug Substance	Dapagliflozin
Study Code	D1692C00012
Edition Number	1
Date	

---

---

**Appendix G**  
**Algorithm on Management of Sustained Elevated Liver Safety**  
**Abnormalities**

---

## **ALGORITHM ON MANAGEMENT OF SUSTAINED ELEVATED LIVER SAFETY ABNORMALITIES**

The monitoring for liver safety will be performed using the serum levels of AST, ALT and TB (see [Figure 1](#) Sustained elevated liver safety abnormalities flow chart).

**Patients with a central laboratory ALT and/or AST >3 X ULN** will be scheduled for a follow-up visit within 3 days following receipt of the initial laboratory results, to obtain repeat central laboratory ALT, AST, TB and Alkaline Phosphatase (ALK-P). In the event that the repeat laboratory assessments cannot be obtained within 3 days, the Investigator is encouraged to discuss possible alternatives with the Sponsor. Patients should remain on study medication until confirmatory results are obtained, unless otherwise contraindicated.

- **If the repeat ALT and AST are  $\leq 3X$  ULN**, patient should continue study treatment according to their original visit schedule unless otherwise contraindicated.
- **If the repeat ALT and/or AST are  $>3X$  ULN but  $\leq 8X$  ULN and TB  $\leq 1.5X$  ULN**, the patient's medical history, including details of risk factors for liver diseases, should be evaluated for potential underlying aetiologies. In addition, specialized blood sampling will be performed to evaluate liver function as well as identify potential causes of laboratory elevation(s). The Investigator should continue to monitor the patient's liver tests every 3 days following receipt of the prior laboratory results until the ALT and AST are  $\leq 2X$  ULN or until ALT and AST are at or below baseline levels. The frequency of retesting can decrease to once a week or less if abnormalities stabilize and the patient is asymptomatic. Patients should remain on study medication unless confirmatory results indicate that a criterion for discontinuation has been met or continuing study medication would be otherwise contraindicated.

**Patients must be discontinued from the study if an initial and repeat confirmatory laboratory tests meet any of the following criteria:**

- ALT and/or AST are  $>3$  x ULN and TB  $>1.5$  x ULN
- ALT and/or AST are  $>5$  x ULN for  $\geq 14$  consecutive days, at any time after initial confirmatory results
- ALT and/or AST are  $>8$  x ULN

In each of these situations, study medication will be discontinued, the Sponsor notified and the End of Treatment Visit performed within 3 days of the confirmed laboratory results (see Section 5.8.2 in the CSP). At the End of Treatment Visit, medical history including details of risk factors for liver diseases (if not previously assessed) will be requested and additional

blood sampling performed (**Specialized Liver Panel and Liver Discontinuation Panel**, see detailed below). Patient should also be scheduled for a Follow-up Visit (ie, procedures of Visit 18) 3 weeks after discontinuation of investigational product. A referral consultation to a hepatologist or gastroenterologist (specializing in liver abnormalities) should be obtained.

Any additional tests and/or examinations should be carried out at the discretion of the Investigator. Any further investigations and laboratory results for patients with abnormal laboratory values at the Follow-up Visit should be made available to the Sponsor upon request.

Additional information, including but not limited to completion of supplemental eCRFs may be requested for certain adverse events and/or laboratory abnormalities which are reported/identified as part of the hepatic safety surveillance.

Following the End of Treatment Visit, the Investigator should continue to monitor the subject's liver tests every 3 days following receipt of the prior laboratory results until the ALT and AST are  $\leq 2 \times$  ULN or until ALT and AST are at or below baseline levels. The frequency of retesting can decrease to once a week or less if abnormalities stabilize and the subject is asymptomatic.

#### **Guidance on Assessment of Hepatic Laboratory Abnormalities**

The following is presented to assist in the evaluation and management of hepatic laboratory values. It is not intended to supplant Investigators' clinical judgment.

Patients who experience ALT and/or AST values  $>3 \times$  ULN confirmed with a repeated test will have the following performed within 3 days of the confirmed laboratory results:

- AE assessment
- Physical Examination for jaundice and other signs of liver diseases
- Review of relevant risk factors and current history focusing on possible causes of the increased ALT and/or AST and/or TB, including:
  - Use of suspect concomitant medication [including over-the-counter (ie, acetaminophen/paracetamol), herbal and vitamin preparations]
  - Recent alcohol consumption or recreational drug/narcotic use
  - Recent unaccustomed physical exertion
  - Occupational or environmental exposure to hepatotoxins
  - Other conditions which may cause liver diseases or which may cause abnormal test results

- Specialized Liver Laboratory Panel (see below)

### **Specialized Liver Panel**

For patients who are being monitored frequently as a result of confirmed AST and/or ALT >3X ULN, additional central laboratory tests will be performed within 3 days of receipt of confirmatory results. These laboratory tests will study the possible causes of the increased ALT and/or AST and/or TB, and may include, but are not limited to, the following tests:

- Hepatitis A IgM
- Hepatitis BsAg
- Hepatitis B Core Ab IgM
- Hepatitis C virus RNA
- Hepatitis C Ab
- Hepatitis E IgM
- Epstein-Barr Virus (EBV) IgM Ab
- Lactate Dehydrogenase (LDH)
- Gamma-glutamyl-transpeptidase (GGT)
- Carbohydrate deficient transferrin (CDT)
- Prothrombin time (PT/INR)
- Iron Panel - iron, ferritin, total iron binding capacity (TIBC)
- Immunology Panel including Antinuclear Antibody (ANA), Anti-Smooth Muscle Antibody (SMA) and Anti-Liver/Kidney Microsomal Antibody (Anti-LKM)
- Anti-tissue Transglutaminase Antibody

### **Liver Discontinuation Panel**


For patients who are discontinued from the study as a result of sustained elevated liver safety abnormalities, additional central laboratory tests will be performed at the time of End of Treatment Visit. Similar to the Specialized Liver Panel, these laboratory tests will study the possible causes of the increased ALT and/or AST and/or TB, and may include, but are not limited to, the following tests:

- Cytomegalovirus (CMV) IgM Ab

- Herpes Simplex Virus (HSV) 1 and 2
- Ceruloplasmin
- Toxoplasmosis
- Alpha-1 antitrypsin

For specific details regarding the Specialized Liver Panel or the Liver Discontinuation Panel laboratory tests, refer to the Central Laboratory Manual for this study.

**Figure 1 Sustained elevated liver safety abnormalities flow chart**



- a In patients with repeat ALT or AST > 3X ULN but ≤ 8X ULN, only patients with TB ≤ 1.5X ULN at Step 1 should be followed according to Step 2b. Patients with an initial TB and confirmatory repeat TB > 1.5X ULN should be followed according to Step 2d.
- b Please see text above in the Appendix for details on additional assessments to be performed (AE assessment, PE, review of current medical history including focused review of risk factors for liver diseases and collection of blood samples [specialized liver panel or liver discontinuation panel]).
- c Confirmatory repeat AST, ALT, ALK-P and TB should be obtained every 3 days following receipt of prior laboratory results, until the ALT and AST are ≤ 2X ULN or until ALT and AST are at or below baseline levels. The frequency of retesting can decrease to once a week or less if abnormalities stabilize and the patient is asymptomatic.



---

**Clinical Study Protocol: Supplement A**

Drug Substance	Dapagliflozin
Study Code	D1692C00012
Supplement Edition Number	1
Supplement Date	

---

---

**Supplement A**  
**Investigators and Study Administrative Structure**

---



Clinical Study Protocol: Supplement A  
Drug Substance Dapagliflozin  
Study Code D1692C00012  
Supplement Edition Number 1  
Supplement Date

## **STAFF AT STUDY SITE(S)**

<b>Centre No.</b>	<b>Centre address</b>	<b>Name (First name, Last name)</b>	<b>Qualifications</b>	<b>Present position</b>	<b>Role in the study</b>
-------------------	-----------------------	---	-----------------------	-------------------------	--------------------------

Clinical Study Protocol: Supplement A  
Drug Substance Dapagliflozin  
Study Code D1692C00012  
Supplement Edition Number 1  
Supplement Date

---

<b>Centre No.</b>	<b>Centre address</b>	<b>Name</b> (First name, Last name)	<b>Qualifications</b>	<b>Present position</b>	<b>Role in the study</b>
-------------------	-----------------------	--	-----------------------	-------------------------	--------------------------

---

Clinical Study Protocol: Supplement A  
Drug Substance Dapagliflozin  
Study Code D1692C00012  
Supplement Edition Number 1  
Supplement Date

---

<b>Centre No.</b>	<b>Centre address</b>	<b>Name</b> (First name, Last name)	<b>Qualifications</b>	<b>Present position</b>	<b>Role in the study</b>
-------------------	-----------------------	--	-----------------------	-------------------------	--------------------------

---

---

<b>Centre No.</b>	<b>Centre address</b>	<b>Name</b> (First name, Last name)	<b>Qualifications</b>	<b>Present position</b>	<b>Role in the study</b>
-------------------	-----------------------	--	-----------------------	-------------------------	--------------------------

---

Clinical Study Protocol: Supplement A  
Drug Substance Dapagliflozin  
Study Code D1692C00012  
Supplement Edition Number 1  
Supplement Date

---

<b>Centre No.</b>	<b>Centre address</b>	<b>Name</b> (First name, Last name)	<b>Qualifications</b>	<b>Present position</b>	<b>Role in the study</b>
-------------------	-----------------------	--	-----------------------	-------------------------	--------------------------

---

---

<b>Centre No.</b>	<b>Centre address</b>	<b>Name</b> (First name, Last name)	<b>Qualifications</b>	<b>Present position</b>	<b>Role in the study</b>
-------------------	-----------------------	--	-----------------------	-------------------------	--------------------------

---

Clinical Study Protocol: Supplement A  
Drug Substance Dapagliflozin  
Study Code D1692C00012  
Supplement Edition Number 1  
Supplement Date

## **ASTRAZENECA STUDY PERSONNEL**

---

<b>Address</b>	<b>Name (First name, Last name)</b>	<b>Qualifications</b>	<b>Present Position</b>	<b>Role in the study</b>
----------------	-------------------------------------	-----------------------	-------------------------	--------------------------

---

Clinical Study Protocol: Supplement A  
Drug Substance Dapagliflozin  
Study Code D1692C00012  
Supplement Edition Number 1  
Supplement Date

---

<b>Address</b>	<b>Name (First name, Last name)</b>	<b>Qualifications</b>	<b>Present Position</b>	<b>Role in the study</b>
----------------	-------------------------------------	-----------------------	-------------------------	--------------------------

---



Clinical Study Protocol: Supplement A  
Drug Substance Dapagliflozin  
Study Code D1692C00012  
Supplement Edition Number 1  
Supplement Date

---

<b>Address</b>	<b>Name (First name, Last name)</b>	<b>Qualifications</b>	<b>Present Position</b>	<b>Role in the study</b>
----------------	-------------------------------------	-----------------------	-------------------------	--------------------------

---

Clinical Study Protocol: Supplement A  
Drug Substance Dapagliflozin  
Study Code D1692C00012  
Supplement Edition Number 1  
Supplement Date

---

<b>Address</b>	<b>Name (First name, Last name)</b>	<b>Qualifications</b>	<b>Present Position</b>	<b>Role in the study</b>
----------------	-------------------------------------	-----------------------	-------------------------	--------------------------

---

Clinical Study Protocol: Supplement A  
Drug Substance Dapagliflozin  
Study Code D1692C00012  
Supplement Edition Number 1  
Supplement Date

---

<b>Address</b>	<b>Name (First name, Last name)</b>	<b>Qualifications</b>	<b>Present Position</b>	<b>Role in the study</b>
----------------	-------------------------------------	-----------------------	-------------------------	--------------------------

---

Clinical Study Protocol: Supplement A  
Drug Substance Dapagliflozin  
Study Code D1692C00012  
Supplement Edition Number 1  
Supplement Date

---

<b>Address</b>	<b>Name (First name, Last name)</b>	<b>Qualifications</b>	<b>Present Position</b>	<b>Role in the study</b>
----------------	-------------------------------------	-----------------------	-------------------------	--------------------------

---

Clinical Study Protocol: Supplement A  
Drug Substance Dapagliflozin  
Study Code D1692C00012  
Supplement Edition Number 1  
Supplement Date

---

<b>Address</b>	<b>Name (First name, Last name)</b>	<b>Qualifications</b>	<b>Present Position</b>	<b>Role in the study</b>
----------------	-------------------------------------	-----------------------	-------------------------	--------------------------

---

Clinical Study Protocol: Supplement A  
Drug Substance Dapagliflozin  
Study Code D1692C00012  
Supplement Edition Number 1  
Supplement Date

---

<b>Address</b>	<b>Name (First name, Last name)</b>	<b>Qualifications</b>	<b>Present Position</b>	<b>Role in the study</b>
----------------	-------------------------------------	-----------------------	-------------------------	--------------------------

---

## DATA MONITORING OR SAFETY COMMITTEE(S)

---

<b>Address</b>	<b>Member name (First name, Last name)</b>	<b>Present position</b>	<b>Role in committee</b>
<b>External Medical Advisor</b>			

---

## **OTHER PARTICIPANTS**

---

<b>Organisation and address</b>	<b>Role in study</b>
---------------------------------	----------------------

---





---

**Clinical Study Protocol Amendment**

Amendment Number	1
Drug Substance	Dapagliflozin
Study Code	D1692C00012
Date	
Protocol Dated	

---

---

**A long term open label study to evaluate the safety and efficacy of dapagliflozin as monotherapy or combination therapies with anti-diabetic drugs in Japanese subjects with type 2 diabetes who have inadequate glycemic control**

---

This submission /document contains trade secrets and confidential commercial information, disclosure of which is prohibited without providing advance notice to AstraZeneca and opportunity to object.

**Sponsor:**

AstraZeneca K.K.

Bristol-Myers K.K.

**Centres affected by the Amendment:**

All centres in the study

**The protocol for the study is to be amended as follows:**

“Previous text” and “Revised text” include only information that has been changed.

**Section of protocol affected:**

Case Identification and Management of Decreased Renal Function, in the Appendix F Case Identification and Management of Decreased Renal Function

**Previous text:**

FOR PATIENTS WHO ARE TREATED WITH METFORMIN:

**If calculated eGFR is <60 mL/min or if there is an increase in serum creatinine of  $\geq 0.5$  mg/dL ( $\geq 44.2$   $\mu\text{mol/L}$ ) above the baseline value based on central laboratory results, this will be considered a case of “decreased renal function”.**

In this circumstance, the investigator should consider evaluating the patient for potentially reversible causes of renal dysfunction including:

1. concurrent use of NSAIDS, antibiotics, or other medications known to affect measures of serum creatinine
2. volume depletion
3. urinary tract infection
4. obstructive uropathy.

Patients should return for repeat central laboratory testing as soon as possible, no later than 7 days after the abnormal result. Investigational drug and metformin should be interrupted pending the results of repeat testing. If after interruption eGFR is still <60 ml/min or if the serum creatinine value remains  $\geq 0.5$  mg/dL ( $\geq 44.2$   $\mu\text{mol/L}$ ) above the baseline value, the patient should permanently discontinue the study medication and be withdrawn from the study (in which case an Adverse Event must be reported).

If after interruption of investigational drug and metformin eGFR is  $\geq 60$  ml/min or if the serum creatinine value has decreased to <0.5 mg/dL (<44.2  $\mu\text{mol/L}$ ) above baseline, investigational drug and metformin can be re-started if appropriate in the judgement of the investigator and following consultation with the study team physician. If the patient restarts investigational drug and metformin and eGFR is again <60 ml/min or the serum creatinine value increases again to  $\geq 0.5$  mg/dL ( $\geq 44.2$   $\mu\text{mol/L}$ ) above baseline, the patient should permanently discontinue the study medication and be withdrawn from the study (in which case an Adverse Event must be reported).

**Revised text:**

FOR PATIENTS WHO ARE TREATED WITH METFORMIN:

**If calculated eGFR is <60 mL/min or if there is an increase in serum creatinine of  $\geq 0.5$  mg/dL ( $\geq 44.2$   $\mu\text{mol/L}$ ) above the baseline value based on central laboratory results, this will be considered a case of “decreased renal function”.**

In this circumstance, the investigator should consider evaluating the patient for potentially reversible causes of renal dysfunction including:

1. concurrent use of NSAIDS, antibiotics, or other medications known to affect measures of serum creatinine
2. volume depletion
3. urinary tract infection
4. obstructive uropathy.

Patients should return for repeat central laboratory testing as soon as possible, no later than 7 days after the abnormal result. Investigational drug and metformin should be interrupted pending the results of repeat testing. If after interruption eGFR is still  $<60$  ml/min or if the serum creatinine value remains  $\geq 0.5$  mg/dL ( $\geq 44.2$   $\mu\text{mol/L}$ ) above the baseline value, the patient should permanently discontinue the study medication and be withdrawn from the study (in which case an Adverse Event must be reported).

If after interruption of investigational drug and metformin the eGFR is  $\geq 60$  ml/min and the serum creatinine value is  $< 0.5$  mg/dL ( $< 44.2$   $\mu\text{mol/L}$ ) above baseline, investigational drug and metformin can be re-started if appropriate in the judgement of the investigator and following consultation with the study team physician.

If the patient restarts investigational drug and metformin and eGFR is again  $< 60$  ml/min or the serum creatinine value increases again to  $\geq 0.5$  mg/dL ( $\geq 44.2$   $\mu\text{mol/L}$ ) above baseline, the patient can then be retested a second time according to the process above. The process for retest can be conducted only twice thorough the study. Thereafter, if the patient restarts investigational drug and metformin and eGFR is again  $< 60$  ml/min or the serum creatinine value increases again to  $\geq 0.5$  mg/dL ( $\geq 44.2$   $\mu\text{mol/L}$ ) above baseline, the patient should permanently discontinue the study medication and be withdrawn from the study (in which case an Adverse Event must be reported).

**Reason for Amendment:**

The change of the process in “Case Identification and Management of Decreased Renal Function” for patients who are treated with metformin.

**Persons who initiated the Amendment:**

Study Delivery Team Leader



---

**Clinical Study Protocol Amendment No 1 Appendix A**

Drug Substance	Dapagliflozin
Study Code	D1692C00012
Edition Number	1
Date	
Protocol Dated	

---

---

**Appendix A**  
**Signatures**

---

## **ASTRAZENECA SIGNATURE(S)**

---

**A long term open label study to evaluate the safety and efficacy of dapagliflozin as monotherapy or combination therapies with anti-diabetic drugs in Japanese subjects with type 2 diabetes who have inadequate glycemc control**

---

This Clinical Study Protocol and all Amendments to the CSP have been subjected to an internal AstraZeneca peer review.

I agree to the terms of this study protocol/amendment.

**AstraZeneca Research and Development  
site representative**

This document contains confidential information, which should not be copied, referred to, released or published without written approval from AstraZeneca. Investigators are cautioned that the information in this protocol may be subject to change and revision.

## **ASTRAZENECA SIGNATURE(S)**

---

**A long term open label study to evaluate the safety and efficacy of dapagliflozin as monotherapy or combination therapies with anti-diabetic drugs in Japanese subjects with type 2 diabetes who have inadequate glycemic control**

---

This Clinical Study Protocol and all Amendments to the CSP have been subjected to an internal AstraZeneca peer review.

I agree to the terms of this study protocol/amendment.

**AstraZeneca Research and Development  
site representative**

This document contains confidential information, which should not be copied, referred to, released or published without written approval from AstraZeneca. Investigators are cautioned that the information in this protocol may be subject to change and revision.

## **ASTRAZENECA SIGNATURE(S)**

---

**A long term open label study to evaluate the safety and efficacy of dapagliflozin as monotherapy or combination therapies with anti-diabetic drugs in Japanese subjects with type 2 diabetes who have inadequate glycemic control**

---

This Clinical Study Protocol and all Amendments to the CSP have been subjected to an internal AstraZeneca peer review.

I agree to the terms of this study protocol/amendment.

**AstraZeneca Research and  
Development site representative**

This document contains confidential information, which should not be copied, referred to, released or published without written approval from AstraZeneca. Investigators are cautioned that the information in this protocol may be subject to change and revision.

Clinical Study Protocol Amendment No 1 Appendix A  
Drug Substance Dapagliflozin  
Study Code D1692C00012  
Edition Number 1  
Date

## **ASTRAZENECA SIGNATURE(S)**

---

**A long term open label study to evaluate the safety and efficacy of dapagliflozin as monotherapy or combination therapies with anti-diabetic drugs in Japanese subjects with type 2 diabetes who have inadequate glycemic control**

---

This Clinical Study Protocol and all Amendments to the CSP have been subjected to an internal AstraZeneca peer review.

I agree to the terms of this study protocol/amendment.

**AstraZeneca Research and  
Development site representative**

This document contains confidential information, which should not be copied, referred to, released or published without written approval from AstraZeneca. Investigators are cautioned that the information in this protocol may be subject to change and revision.





---

**Clinical Study Protocol Amendment**

Amendment Number	2
Drug Substance	Dapagliflozin
Study Code	D1692C00012
Date	
Protocol Dated	

---

---

**A long term open label study to evaluate the safety and efficacy of dapagliflozin as monotherapy or combination therapies with anti-diabetic drugs in Japanese subjects with type 2 diabetes who have inadequate glycemic control**

---

This submission /document contains trade secrets and confidential commercial information, disclosure of which is prohibited without providing advance notice to AstraZeneca and opportunity to object.

**Sponsor:**

AstraZeneca K.K.

Bristol-Myers K.K.

**Centres affected by the Amendment:**

All centres in the study

**The protocol for the study is to be amended as follows:**

“Previous text” and “Revised text” include only information that has been changed.

## **Change 1**

### **Section of protocol affected:**

7.1 Volume of blood (Last paragraph)

### **Previous text:**

None

### **Revised text:**

Some of the blood collection tubes and related devices (hereinafter referred to as clinical laboratory test kits) that will be provided by the sponsor have not been approved yet in Japan. In addition, devices already approved in Japan that are imported from overseas and provided by the sponsor are treated as unapproved clinical laboratory test kits. The provided BD Vacutainer<sup>®</sup> Holders, BD Vacutainer<sup>®</sup> SST<sup>™</sup> Tubes, BD Vacutainer<sup>®</sup> EDTA Tubes, BD Vacutainer<sup>®</sup> Citrate Tubes, BD Vacutainer<sup>®</sup> Safety-Lok<sup>™</sup> Blood Collection Sets, BD Vacutainer<sup>®</sup> Blood Collection Needles and 2 mL BD Vacutainer<sup>®</sup> Fluoride Tubes are unapproved medical devices in Japan. Further details of the clinical laboratory test kits used in this study are described in Supplement B.

During the study period (from the start of screening period to the completion of follow-up period), any failure of the unapproved clinical laboratory test kits must be reported to the sponsor. See Section 13.4 for handling of any failure associated with serious adverse events.

Used/un-used clinical laboratory test kits will be appropriately destroyed at the study site.

### **Reason for Amendment:**

Due to the partial amendment to the handling of notifications of plans for clinical trials related to machinery and equipment etc.

## **Change 2**

### **Section of protocol affected:**

8.1 Ethical conduct of the study

### **Previous text:**

The study will be performed in accordance with ethical principles that have their origin in the Declaration of Helsinki and are consistent with ICH/Good Clinical Practice, applicable regulatory requirements and the AstraZeneca policy on Bioethics and Human Biological Samples.

The applicable regulatory requirements in Japan are ‘Good Clinical Practice for Trials on Drugs (MHLW Ordinance No. 28, 27 March 1997, partially revised by MHLW Ordinance and their related notifications).

**Revised text:**

The study will be performed in accordance with ethical principles that have their origin in the Declaration of Helsinki and are consistent with ICH/Good Clinical Practice, applicable regulatory requirements and the AstraZeneca policy on Bioethics and Human Biological Samples.

The applicable regulatory requirements in Japan are ‘Good Clinical Practice for Trials on Drugs (MHLW Ordinance No. 28, 27 March 1997, partially revised by MHLW Ordinance and their related notifications) and GCP for Trials on Medical Devices (MHLW Ordinance No. 36 issued in 2005, partially revised by MHLW Ordinance and their related notifications).

**Reason for Amendment:**

Due to the partial amendment to the handling of notifications of plans for clinical trials related to machinery and equipment etc.

**Change 3**

**Section of protocol affected:**

13.4 Report on Medical Device Failures

**Previous text:**

None

**Revised text:**

Failure of BD Vacutainer® Holders, BD Vacutainer® SST™ Tubes, BD Vacutainer® EDTA Tubes, BD Vacutainer® Citrate Tubes, BD Vacutainer® Safety-Lok™ Blood Collection Sets, BD Vacutainer® Blood Collection Needles and 2 mL BD Vacutainer® Fluoride Tubes provided by the sponsor (see Supplement B) (failures in a broad sense including breakage or glitches of devices) which may be the cause of certain serious adverse events, judged by the investigators, should be reported to the study sponsor along with the serious adverse events within 1 day.

**Reason for Amendment:**

Due to the partial amendment to the handling of notifications of plans for clinical trials related to machinery and equipment etc.

## Change 4

### Section of protocol affected:

Supplement B Key features of clinical laboratory test kits

### Previous text:

None

### Revised text:

#### 1. KEY FEATURES

Key features of the kits with identifier K56C1-001, K56C2-001, K56C2-006 (3 mL/6 mL), K56C2-008, K56C2-019, K56C2-031 and K56C2-033 are shown in [Table 1](#) and [Table 2](#).

**Table 1** Key features of K56C2-001, K56C2-006 (6 mL), K56C2-008, K56C2-019 and K56C2-031 (Approved in Japan)

<u>Item</u>	<u>Key feature</u>				
<u>Identifier</u>	<u>K56C2-001</u>	<u>K56C2-006 (6 mL)</u>	<u>K56C2-008</u>	<u>K56C2-019</u>	<u>K56C2-031</u>
<u>Overseas brand name</u>	<u>BD Vacutainer<sup>®</sup> SST<sup>™</sup> Tubes</u>	<u>BD Vacutainer<sup>®</sup> EDTA Tubes</u>	<u>BD Vacutainer<sup>®</sup> Citrate Tubes</u>	<u>BD Vacutainer<sup>®</sup> Safety-Lok<sup>™</sup> Blood Collection Sets</u>	<u>BD Vacutainer<sup>®</sup> Blood Collection Needles</u>
<u>Manufacturers</u>	<u>Becton, Dickinson And Company</u>	<u>Becton, Dickinson And Company</u>	<u>Becton, Dickinson And Company</u>	<u>Becton, Dickinson And Company</u>	<u>Becton, Dickinson And Company</u>
<u>Categorisation</u>	<u>56</u>	<u>56</u>	<u>56</u>	<u>56</u>	<u>56</u>
<u>Planned generic name</u>	<u>Blood Collection Tubes</u>	<u>Blood Collection Tubes</u>	<u>Blood Collection Tubes</u>	<u>Blood Collection Needles</u>	<u>Blood Collection Needles</u>
<u>Planned classification</u>	<u>2</u>	<u>2</u>	<u>2</u>	<u>2</u>	<u>2</u>

**Table 1 Key features of K56C2-001, K56C2-006 (6 mL), K56C2-008, K56C2-019 and K56C2-031 (Approved in Japan)**

<u>Item</u>	<u>Key feature</u>				
<u>Identifier</u>	<u>K56C2-001</u>	<u>K56C2-006 (6 mL)</u>	<u>K56C2-008</u>	<u>K56C2-019</u>	<u>K56C2-031</u>
<u>Package insert</u>	<u>BD Vacutainer® Blood Collection Tubes (13 mm SST blood collection tube, Draw volume 3.5 mL/5 mL/8.5 mL)</u>	<u>BD Vacutainer® Blood Collection Tubes (EDTA blood collection tube, Hemogard, Draw volume 6 mL)</u>	<u>BD Vacutainer® Blood Collection Tubes (Sodium Citrate blood collection tube, Hemogard, Draw volume 4.5 mL)</u>	<u>BD Vacutainer® Blood Collection Sets (Safety-Lock)</u>	<u>BD Vacutainer® System (Multi-sample needles)</u>

**Table 2 Key features of K56C1-001, K56C2-006 (3 mL) and K56C2-033 (Similar products are approved in Japan)**

<u>Item</u>	<u>Key feature</u>		
<u>Identifier</u>	<u>K56C1-001</u>	<u>K56C2-006 (3 mL)</u>	<u>K56C2-033</u>
<u>Overseas brand name</u>	<u>BD Vacutainer® Holders</u>	<u>BD Vacutainer® EDTA Tubes</u>	<u>2 mL BD Vacutainer® Fluoride Tubes</u>
<u>Manufacturers</u>	<u>Becton, Dickinson And Company</u>	<u>Becton, Dickinson And Company</u>	<u>Becton, Dickinson And Company</u>
<u>Categorisation</u>	<u>56</u>	<u>56</u>	<u>56</u>
<u>Planned generic name</u>	<u>Vacutainer Holder</u>	<u>Blood Collection Tubes</u>	<u>Blood Collection Tubes</u>
<u>Planned classification</u>	<u>1</u>	<u>2</u>	<u>2</u>
<u>Package insert</u>	<u>BD Vacutainer® System (BD Vacutainer® single-use holder plus)</u>	<u>BD Vacutainer® Blood Collection Tubes (EDTA blood collection tube, Hemogard, Draw volume 3 mL)</u>	<u>BD Vacutainer® Blood Collection Tubes (13 mm Sodium fluoride/EDTA 2 Na blood collection tube, Hemogard, Draw volume 2 mL)</u>

## 2. OTHER KEY FEATURES

Regarding devices with the identifiers K56C2-001, K56C2-006 (6 mL), K56C2-008, K56C2-019 and K56C2-031, for other features, shapes, structures, materials, intended use, item specification, direction for use, and precaution for use, see the package insert “BD Vacutainer® Blood Collection Tubes”, “BD Vacutainer® Blood Collection Sets” and “BD Vacutainer® System” presented in Table 1.

Devices with the identifiers K56C1-001, K56C2-006 (3 mL) and K56C2-033, to be used in this study are Vacutainer Holder (Reusable type), 13 mm Blood Collection Tube (EDTA K2, Hemogard, Draw volume 3 mL and 13 mm Blood Collection Tube (Fluoride and Oxalate, Hemogard, Draw volume 2 mL). There are similar devices which have already been approved in Japan. For other features, shapes, structures, materials, intended use, item specification, direction for use and precaution for use, see the package insert of “BD Vacutainer System (BD Vacutainer Single Use Holder Plus)” and “BD Vacutainer® Blood Collection Tubes” presented in Table 2. The differences between the unapproved medical device used in this study and the similar medical device approved in Japan are presented in Table 3.

**Table 3** **Difference of medical device**

<u>Identifier</u>	<u>Unapproved medical device used in this study</u>	<u>Similar medical device approved in Japan</u>	<u>Differences</u>
<u>K56C1-001</u>	<u>Vacutainer Holder (Yellow, Reusable type)</u>	<u>Vacutainer Holder (White, Not reusable type)</u>	<u>Colour and reusable</u>
<u>K56C2-006 (3 mL)</u>	<u>13 mm Blood Collection Tube (EDTA K2, Hemogard, Draw volume 3 mL, Paper label)</u>	<u>13 mm Blood Collection Tube (EDTA K2, Hemogard, Draw volume 3 mL, See thru label)</u>	<u>Label</u>
<u>K56C2-033</u>	<u>13 mm Blood Collection Tube (Fluoride and Oxalate, Hemogard, Draw volume 2 mL)</u>	<u>13 mm Blood Collection Tube (Acid Citric Dextrose, Hemogard, Draw volume 2 mL)</u>	<u>Type of additive</u>

### **Reason for Amendment:**

Due to the partial amendment to the handling of notifications of plans for clinical trials related to machinery and equipment etc.

### **Persons who initiated the Amendment:**

Study Leader

## 機械器具 56 採血又は輸血用器具

管理医療機器 真空密封型採血管 35414000

## BD バキュテイナ採血管


## 再使用禁止

## 【禁忌・禁止】

- 再使用禁止
- 本品を使用して採取した血液を体内に戻さないこと。[採取した血液の安全性が確保できない。]
- 本品を使用し採血する際には、耐圧性能を有するゴムスリーブ付き採血針と単回使用採血ホルダー（以下「ホルダー」と呼ぶ）との組み合わせ以外では使用しないこと。
- 採血管が室内温度に戻らないうちに採血を行わないこと。[採血管の温度により採血管内の圧力が変化し、採血管内の内容物等が患者の体内に逆流するおそれがある。]
- 採血管を抜くまで、患者の腕の血管の圧迫を解除したり、動かしたりしないこと。[圧迫を解除した際、あるいは腕の配置によっては静脈血圧が急激に低下し、採血管内の内容物等が患者の体内に逆流するおそれがある。]
- 採血管に血液が流入し始めた後は、ホルダーに押し込むような力を採血管に加えないこと。[採血管内の圧力が変化し、採血管内の内容物等が患者の体内に逆流するおそれがある。]
- 採血終了後、採血管に採血針が刺さったままの状態で駆血帯を外さないこと。[駆血帯を外すことによる圧力の変動により、採血管内の内容物等が患者の体内に逆流するおそれがある。]
- ホルダーは患者ごとの使用とし、使用後は廃棄すること。[ホルダーに血液が付着した場合は、交差感染のおそれがある。]
- 体外循環回路又は中心静脈から採血は行わないこと。[圧力の変動により、採血管内の内容物等が患者の体内に逆流するおそれがある。]

## 【形状、構造及び原理等】

- 本品は、プラスチック製又はガラス製で、採血管内部に採血量分の陰圧がかけられた真空採血管である。
- キャップの形状はコンベンショナルタイプ（ゴム栓タイプ）とヘモガードタイプ（ゴム栓とプラスチックカバーが一体になったキャップ）がある。



採血管サイズ・採血量は箱のラベルを参照のこと。

## 【使用目的、効能又は効果】

本品は、各種血液検査のために血液を採取する滅菌済み真空採血管である。

## 【品目仕様等】

JIS T3233 真空採血管に適合する。

## 【操作方法又は使用方法等】

1. 室内温度になった本品を準備する。
2. あらかじめ手袋を着用する。
3. 採血針をホルダーにセットする。
4. 駆血帯を装着し、穿刺部位を消毒する。
5. 採血針を血管に穿刺し、本品をホルダーにまっすぐ完全に押し込む。
6. 採血の血流が停止したら、直ちに本品をホルダーから外す。
7. 連続採血する場合には、ホルダーを固定したまま、本品を取り替える。
8. 採血終了後、本品をホルダーから抜去した後に駆血帯を外す。
9. 抜針し、採血針をホルダーから外さずに適切な耐貫通性廃棄容器に廃棄する。

## 〈使用方法に関連する使用上の注意〉

- 採血後直ちにゆっくりと必要回数転倒混和すること。[転倒混和が不十分な場合、抗凝固剤入り品種では血液凝固が発生する可能性があり、また凝固促進剤入り品種では凝固不十分となる場合がある。]

## 〈転倒混和回数〉

採血管の種類	回数
プラスチック製プレイン採血管	5
SST/SST II 採血管 (血清分離剤入り)	5
ヘパリン入り採血管	8～10
PST 採血管 (ヘパリンリチウム/血漿分離剤入り)	8～10
フッ化ナトリウム入り採血管	8～10
EDTA 入り採血管	8～10
クエン酸ナトリウム緩衝液入り採血管	3～4

- 遠心分離機によって血清を得る品種では、転倒混和後凝固時間を十分にとってから遠心分離を行うこと。[遠心分離後フィブリンが析出したり、SST/SST II 採血管（血清分離剤入り）は血清側の管壁に血清分離剤が付着する可能性がある。]

## 〈凝固時間〉

採血管の種類	時間(分)
プレイン採血管	60
SST/SST II 採血管 (血清分離剤入り)	30
トロンビン入り採血管	5

- 遠心条件は以下を推奨する。[遠心強度が足りない場合や遠心時間が短い場合は、血液が十分に分離せず、正確な検査値が得られない可能性がある。]

〈推奨する遠心条件〉

採血管の種類	RCF (g)	時間(分)
13mm SST 採血管 (血清分離剤入り) PST 採血管 (ヘパリンリチウム / 血漿分離剤入り)	1100-1300	10
16mm SST 採血管 (血清分離剤入り)	1000-1300	10
SST II 採血管 (血清分離剤入り)	1300-2000	10
ヘパリン入り採血管 フッ化ナトリウム入り採血管 EDTA 入り採血管	1300 以下	10
クエン酸ナトリウム緩衝液 入り採血管	1500	15
PPT 採血管 (EDTA2K/ 血漿分離剤入り)	1100	10
CPT 単核球分離用採血管 (ヘパリンナトリウム入り)	1500-1800	15
CPT 単核球分離用採血管 (クエン酸ナトリウム緩衝液入り)		20


- SST/SST II 採血管 (血清分離剤入り) は 2 回以上遠心分離しないこと。[検査値に影響を及ぼす可能性がある。]
- SST/SST II 採血管 (血清分離剤入り) を遠心分離後、凍結保存する場合は、血清を別容器に移し替えること。[採血管のまま凍結した場合、解凍時、血球内成分が血清中に移動し、検査値に影響を及ぼす可能性がある。]
- エアシューターを使用する際はキャップ部分が重ならないようにし、クッション材等を用いて採血管を保護してから使用すること。[キャップ部分が重なった状態で衝撃を受けると、キャップが破損したり外れたりする可能性がある。]
- \*本品を使用し採血する際には、BD バキュテイナシステムのホルダー及び採血針/ルアーアダプター、BD バキュテイナブラッドコレクションセット (医療機器認証番号 220AFBZX00064000)、セーフティーロック・ウイングコレクションセット (医療機器認証番号 16200BZZ00144000)、ウイング・コレクション・セット (医療機器認証番号 16200BZZ00144000) の組合せが望ましい。

## 【使用上の注意】

### 〈重要な基本的注意〉

- 患者の腕及び採血管の底部が採血中常に下向きであることを確認すること。
- 翼状針を使用して採血する際は、採血管の位置が上下に動かないようにすること。[採血管内圧と静脈圧の関係から採血管内の内容物が患者の体内に逆流するおそれがある。]
- 採血時及び採血管を取り扱う際は、血液との接触を最小限にするため、手袋をする等の適切な標準予防策をとること。
- 採血後、直ちにゆっくりと転倒混和し、血液と添加剤を十分に混和させること。
- 使用後の採血管は適切な廃棄容器に捨てること。
- 血清採取用の採血管を使用する場合は、採取した血液を十分に凝固させた後、血清を採取すること。
- 採取した血液が飛散する事があるため、栓を親指で押し上げて外さないこと。ヘモガードタイプの栓は下記の方法で外すこと。
  1. 栓の下に親指をあてて、片方の手で採血管を握る。親指で栓を押し上げながら、もう一方の手で栓がゆるむところまで回す。(図 1)

2. 栓を持ち上げる前に、親指を栓から離す。(図 2)
3. 栓を持ち上げて採血管から外す。プラスチックカバーがゴム栓から外れてしまった場合には、再度プラスチックカバーをゴム栓に装着せず、ゴム栓を採血管から注意深く取り外す。(図 3)



- 破損や異物の混入が認められる採血管は使用しないこと。
- 針刺し損傷等によって血液曝露した場合は、直ちに流水でよく洗い、施設のプロトコールに従い、適切な医療処置を受けること。
- 原則として、SST/SST II 採血管 (血清分離剤入り) は生化学検査、免疫血清学検査、また PST 採血管 (ヘパリンリチウム / 血漿分離剤入り) は生化学検査に用いること。
- 薬物血中濃度の測定を行う場合には適切な採血管を使用すること。[検査値に影響を及ぼすおそれがある。]
- \*他社ホルダーと併用する場合には、他社ホルダーの添付文書の注意事項を確認の上、使用すること。[16mm ヘモガードタイプ採血管 (キャップ外径: 17.5mm) を他社ホルダーと併用した場合、採血管をホルダーから抜く際にヘモガードキャップが抜ける可能性がある。]
- 使用期限の切れた採血管は使用しないこと。
- ガラス採血管は破損する可能性があるため、操作には十分注意すること。
- 血液の入ったガラス採血管を 0℃以下に保存すると、破損するおそれがあるため避けること。
- 再栓するときは、採血管に栓をかぶせ、ゴム栓がしっかりと採血管に入るまで、回しながら押し込むこと。
- 採血管をそのまま用いて検査する場合は、分析装置メーカーに適合の可否を確認し、必要に応じて調整してから使用すること。
- 使用期限内でもプラスチック管内部に徐々に空気が流入し、採血量が減衰する可能性がある。
- 採血時の気圧、血圧、温度等の影響により、採血量が変化する可能性がある。
- ゲル内に泡が認められる場合があるが、検査値への影響はない。

## 【貯蔵・保管方法及び使用期間等】

### 〈貯蔵・保管方法〉

水ぬれに注意し、直射日光を避け、4~25℃ (CPT 採血管のみ 18~25℃) で保管すること。

### 〈使用期限〉

使用期限は採血管に印字又は採血管貼付ラベルに表示 [自己認証「社内データ」による。]

## 【包装】

- 1 箱 60 本入り
- 1 箱 100 本入り
- 1 箱 1000 本入り

## 【製造販売業者及び製造業者の氏名又は名称及び住所等】

製造販売業者:

日本ベクトン・ディッキンソン株式会社  
〒960-2152 福島県福島市土船字五反田1番地  
TEL: 0120-8555-90 (BDお客様情報センター)

外国製造業者:

ベクトン・ディッキンソンアンドカンパニー  
(Becton, Dickinson and Company)  
国名: アメリカ合衆国、イギリス





## 機械器具 56 採血又は輸血用器具

管理医療機器 単回使用採血用針 35209002

## BD バキュテイナ ブラッドコレクションセット

## 再使用禁止


## 【禁忌・禁止】

- 再使用禁止
- 真空採血管とホルダーの組み合わせで採血する場合、滅菌済み真空採血管および単回使用採血ホルダーとの組み合わせ以外では使用しないこと。
- 採血終了後、採血管に採血針が刺さったままの状態で駆血帯を外さないこと。[駆血帯を外すことによる圧力の変動により、採血管内の内容物等が患者の体内に逆流するおそれがある。]
- ホルダーは患者ごとの使用とし、使用後は廃棄すること。[ホルダーに血液が付着した場合は、交差感染のおそれがある。]


## 【形状・構造及び原理等】

本品はチューブの一端にルーアコネクタ及びルーアダプターが、他端には翼付針が接続されている。使用後の針部分を保護、収納するロック機構を備えている。

## 1. セーフティロック・ブラッドコレクションセット



## 2. プッシュボタン・ブラッドコレクションセット



本品は延長チューブ部分にポリ塩化ビニル（可塑剤：フタル酸ジ(2-エチルヘキシル)）を使用。

## 【使用目的、効能又は効果】

本品は血液検査のため、静脈に穿刺し、真空採血管を用いた血液検体の採取に用いる、ディスポーザブル翼付針である。

## 【品目仕様等】

JIS T3220 の下記項目に適合している。

外観及び清浄度	引抜き強さ
寸法の許容差	気密性
潤滑剤の量	引張試験
外観及び清浄度(採血針の内面)	導管
ネクタ及びルーアダプターのはめあい	ゴムスリーブの耐圧性
刃先	ホルダー(ホルダー付のみ)


## 【操作方法又は使用方法等】

## 1. セーフティロック・ブラッドコレクションセットの場合

- 1) あらかじめ手袋を着用する。
- 2) 開封して本品を取り出す。
- 3) 駆血帯を装着し、穿刺部位を消毒する。

- 4) 針カバーをまっすぐ引いて取り外し（図1）、図のように翼部（ウイング部）を持ち穿刺する（図2）。

注意：セーフティシールドを持って穿刺しないこと。




- 5) ホルダーに採血管をまっすぐ押し込む。
- 6) 採血の血流が停止したら、直ちに採血管をホルダーから外す。
- 7) 連続採血する場合には、ホルダーを固定したまま採血管を取り替える。
- 8) 採血終了後、採血管をホルダーから除去した後に駆血帯を外す。
- 9) 抜針し、ホルダーごと適切な耐貫通性廃棄容器に廃棄する。

## 〈抜針方法〉

- 採血終了後、次のいずれかの方法で抜針する。穿刺部位には滅菌ガーゼをあてがう。

- 1) 親指と人差し指でセーフティシールドのグリップ部分とチューブをつまんで抜針する（図3）。
- 2) どちらか一方の翼部（ウイング部）あるいは両方の翼部（ウイング部）をつまんで抜針する。（図4）。



## 〈セーフティシールドの使用方法〉

- 1) 片手操作1（図5）：

抜針後、片手でチューブを持ち、親指と人差し指を使ってカチッと音がするまでセーフティシールドを前方に動かす。針は完全にセーフティシールド内に収納され、同時にセーフティシールドはロックされる（図8）。

- 2) 片手操作2（図6）：

滅菌ガーゼをあてがっている方の手の親指と人差し指でチューブをつまみ、カチッと音がするまでセーフティシールドを前方に動かす（図6）。針は完全にセーフティシールド内に収納され、同時にセーフティシールドはロックされる（図8）。

- 3) 両手操作（図7）：


片手のどちらか一方の翼部（ウイング部）をつまみ、もう一方の手でセーフティシールドのグリップ部分をつまむ。カチッと音がするまで翼部（ウイング部）をセーフティシールドの後部にスライドさせる。針は完全にセーフティシールド内に収納され、同時にセーフティシールドはロックされる（図8）。

注意：セーフティシールドの胴体部分をつままないこと。

[セーフティシールドの作動が妨害されることがある。]

注意：セーフティシールドに過度の力を加えないこと。

[セーフティシールドが破損するおそれがある。]




2. プッシュボタン・ブラッドコレクションセットの場合

- 1) あらかじめ手袋を着用する。
- 2) 開封して本品を取り出す。
- 3) ルアーアダプターをホルダーにセットする。
- 4) 駆血帯を装着し、穿刺部位を消毒する。
- 5) 針カバーをまっすぐ引いて取り外し、次のいずれかの方法で穿刺する。


●親指と人差し指で翼部（ウイング部）を持ち穿刺する（図9）。

●本体部分を持ち穿刺する（図10）。


注意：本体部分を持って穿刺する場合、誤ってプッシュボタンを押して安全装置を作動させないように注意すること。




- 6) 血液のバックフローを確認する（図11）。



- 7) 採血後、採血管をホルダーから外した後に駆血帯を外し、滅菌ガーゼを穿刺部位にあてがいが針管の先端を覆う。穿刺したままの状態では本体を持ち、プッシュボタン（図12）を作動させる（図13）。



注意：安全装置作動の妨げになるため、本体部分後部のチューブを触らないこと（図14）。



- 8) 針が完全に本体に収納されたことを確認する。

注意：安全装置が作動しなかった場合は、もう一度プッシュボタンを押すこと。それでも作動しない場合は、ゆっくりと抜針し、速やかに耐貫通性廃棄容器に廃棄すること。

- 9) 適切な耐貫通性廃棄容器に廃棄する。

〈使用方法に関連する使用上の注意〉

- 採血時及び採血管を取り扱う際は、血液との接触を最小限にするため、手袋をするなどの適切な標準予防策をとること。
- 消毒後は、穿刺部位を触らないこと。

- 採血針をホルダーに確実に固定すること。[ホルダーへの固定が確実でない場合、採血針が使用中に外れるおそれがある。]
- 針カバーを外す際は、針先が針カバーに触れないように注意すること。[刃先が変形するおそれがある。]
- ゴム栓タイプ以外の採血管をホルダーに挿入する際、採血管をゆっくりとまっすぐ挿入し最後まで完全に押し込まないこと。[ゴムスリーブの外れなどにより血液曝露が起こるおそれがある。]
- 1本のルアーアダプターから採血する採血管本数は原則として6本までとすること。[ゴムスリーブから血液漏れが起こるおそれがある。]  
※日本臨床検査標準協議会（JCCLS）標準採血法ガイドライン参照。
- ホルダー内に血液漏れが生じた場合は、採血針を直ちに血管から抜去し、新しいルアーアダプター、採血針及び専用ホルダーと交換すること。[採血管を多数使用した場合、ゴムスリーブが戻らず、血液漏れが起こるおそれがある。]
- 安全装置が正常に作動しなかった場合や、作動時に破損した場合は、針刺し損傷に注意し速やかに耐貫通性廃棄容器に廃棄すること。

【使用上の注意】

〈重要な基本的注意〉

- 包装が破損・汚染している場合や、製品に破損等の異常が認められる場合は使用しないこと。
- 包装を開封したらすぐに使用し、使用後は感染防止に留意して安全な方法で処分すること。
- あらかじめ各接合部に緩みがないことを確認してから使用すること。また、使用中は定期的に緩み、外れ、破損、血液漏れがないことを確認すること。
- 採血針及びゴムスリーブ（ルアーアダプター）に直接手を触れないこと。[針刺し損傷のおそれがある。]
- 針刺し損傷が起きた場合は、施設のプロトコルに従い、直ちに適切な処置をすること。
- 本品のルアーコネクタは国際規格のルアーフィッティングで規定されている規格に準拠しているが、接続相手が同様の規格に準拠している場合でも締め方や接続部の取扱い等により、接続が緩む場合が想定される。確実にしっかりと接続し、使用中は本品の破損、接続部のゆるみ及び血液漏れ等について確認すること。
- 採血中に採血針がホルダーから外れた場合、直ちに患者から抜針し、血液との接触を避け、感染防止措置を取ること。
- チューブを採血針の先端、刃物その他鋭利物等で傷をつけないよう注意すること。
- 各接合部に過度の負荷を加えないこと。[破損のおそれがある。]
- 使用後の製品は適切な耐貫通性廃棄容器に廃棄すること。

【貯蔵・保管方法及び使用期間等】

〈貯蔵・保管方法〉

水ぬれに注意し、直射日光及び高温・多湿を避けて保管すること。

〈使用期限〉

箱に記載。

【包装】

1箱 25個入り

1箱 50個入り

【製造販売業者及び製造業者の氏名又は名称及び住所等】

製造販売業者：

日本ベクトン・ディッキンソン株式会社

〒960-2152 福島県福島市土船字五反田1番地

TEL：0120-8555-90（BDお客様情報センター）

外国製造業者：

ベクトン・ディッキンソン アンド カンパニー

（Becton, Dickinson and Company）

国名：アメリカ合衆国



## 機械器具 56 採血又は輸血用器具

管理医療機器 単回使用採血用針 35209002

## BD バキュテイナシステム

## 再使用禁止


## 【禁忌・禁止】

- 再使用禁止
- 採血針とホルダーの組合せで採血する場合、耐圧性能を有するゴムスリーブ付採血針と単回使用採血ホルダーの組み合わせ以外では使用しないこと。
- 採血終了後、採血管に採血針が刺さったままの状態で駆血帯を外さないこと。[駆血帯を外すことによる圧力の変動により、採血管内の内容物等が患者の体内に逆流するおそれがある。]
- ホルダーは患者ごとの使用とし、使用後は廃棄すること。[ホルダーに血液が付着した場合は、交差感染のおそれがある。]


## \*【形状・構造及び原理等】

本品には下記のような種類がある。

1. マルチ採血針
2. フラッシュバックニードル（血液のバックフロー確認機能付）
3. エクリプス採血針
4. ホルダー付エクリプス採血針
5. ルアーアダプター（翼状針等を接続する為のアダプター）
6. シングルユースホルダー（単回使用採血ホルダー）
7. ブラッド・トランスファー・デバイス（血液分注器具）




## 5. ルアーアダプター



## 6. シングルユースホルダー



## 7. ブラッド・トランスファー・デバイス



針（使用されている場合）の材質：ステンレス

本品（シングルユースホルダーを除く）に使用されているゴムスリーブは、耐圧性能を有している。

## 【使用目的、効能又は効果】

血液検査のため、静脈に穿刺し、真空採血管を用いた血液検体の採取に用いる。

## 【品目仕様等】

JIS T3220 の下記項目に適合している。

外観及び清浄度	ゴムスリーブの耐圧性
寸法の許容差	ホルダー（ホルダー付のみ）
潤滑剤の量	
外観及び清浄度	
刃先	
引き抜き強さ	
弾性	
曲げ強さ	

## 【操作方法又は使用方法等】






## (1) マルチ採血針、フラッシュバックニードルの場合

- 1) あらかじめ手袋を着用する。
- 2) キャップを回転させた後、まっすぐ引き抜いて外す。
- 3) 採血針をホルダーにセットする。
- 4) 駆血帯を装着し、穿刺部位を消毒する。
- 5) 針カバーをまっすぐ引き抜いて外し、採血針を血管に穿刺する。
- 6) 採血管をホルダーにまっすぐ押し込む。
- 7) 採血の血流が停止したら、直ちに採血管をホルダーからまっすぐ外す。
- 8) 連続採血する場合には、ホルダーを固定したまま、採血管を取り替える。
- 9) 採血終了後、採血管をホルダーから除去した後に駆血帯を外す。
- 10) 抜針し、採血針をホルダーから外さず適切な耐貫通性廃棄容

器に廃棄する。

## \* (2) エクリプス採血針、ホルダー付エクリプス採血針の場合

- 1) あらかじめ手袋を着用する。
- 2) 安全シールドと針カバーの両方をもち、キャップを回転させた後(図1)、まっすぐ引いて外す。
- 3) 採血針をホルダーにセットする(ホルダー付エクリプス採血針の場合は不要)。
- 4) 駆血帯を装着し、穿刺部位を消毒する。
- 5) 安全シールドをホルダー側にそって引き(図2)、針カバーをまっすぐ引き抜いて外し(図3)、採血針を血管に穿刺する。
- 6) 採血管をホルダーにまっすぐ押し込む。
- 7) 採血の血流が停止したら、直ちに採血管をホルダーからまっすぐ外す。
- 8) 連続採血する場合には、ホルダーを固定したまま、採血管を取り替える。
- 9) 採血終了後、採血管をホルダーから除去した後に駆血帯を外す。
- 10) 抜針後、ホルダーを保持している手の親指を、親指パッド部に当て(図4)、クリック音が鳴るまで安全シールドを作動させる(図5)。  
注意：親指パッド以外を用いて作動させないこと。[針刺し損傷のおそれがある。]  
注意：机などで安全シールドを作動させないこと。[安全シールドの破損や血液飛散のおそれがある。]
- 11) 採血針をホルダーから外さず適切な耐貫通性廃棄容器に廃棄する。




## (3) ブラッド・トランスファー・デバイスの場合

- 1) 採血後、採血針等を外したシリンジの先にブラッド・トランスファー・デバイスを確実に取り付ける。  
注意：シリンジが確実に接続されていない場合、血液が漏れ出るおそれがある。
- 2) シリンジの先端を下向きにし、ブラッド・トランスファー・デバイスに採血管または血液培養ボトルを押し込む。
- 3) 分注終了後、シリンジとブラッド・トランスファー・デバイスを取り付けたまま廃棄する。

### <使用方法に関連する使用上の注意>

- 採血時及び採血管を取り扱う際は、血液との接触を最小限にするため、手袋をするなどの適切な標準予防策をとること。
- 消毒後は、穿刺部位を触らないこと。
- キャップを外す際は、キャップがゴムスリーブに触れないよう注意すること。[下図のようにゴムスリーブから後方針が飛び出し、採血時に血液漏れが起こるおそれがある。]



- 採血針をホルダーに確実に固定すること。[ホルダーへの固定が確実でない場合、採血針が使用中に外れるおそれがある。]
- 安全シールドをねじったり過度な力を加えたりしないこと。

[破損するおそれがある。]

- 針カバーを外す際は、針先が針カバーに触れないように注意すること。[刃先が変形するおそれがある。]
- ゴム栓タイプ以外の採血管をホルダーに挿入する際、採血管をゆっくりとまっすぐ挿入し最後まで完全に押し込まないこと。[ゴムスリーブの外れなどにより血液曝露が起こるおそれがある。]
- 1本の採血針により採血する採血管本数は原則として6本までとすること。[ゴムスリーブから血液漏れが起こるおそれがある。]  
※日本臨床検査標準協議会(JCCLS)標準採血法ガイドライン参照。
- ホルダー内に血液漏れが生じた場合は、採血針を直ちに血管から抜去し、新しいルアーアダプター、採血針及び専用ホルダーと交換すること。[採血管を多数使用した場合、ゴムスリーブが戻らず、血液漏れが起こるおそれがある。]
- 安全シールドが正常に作動しなかった場合や、作動時に破損した場合は、針刺し損傷に注意し速やかに耐貫通性廃棄容器に廃棄すること。

### <使用上の注意>

#### <重要な基本的注意>

- 針カバーを外したら、直ちに使用すること。また、使用後は針刺し損傷を防止するために使用方法に従い、感染防止に留意し、耐貫通性廃棄容器に廃棄すること。
- 包装が破損・汚染している場合や製品に破損等の異常が認められる場合は使用しないこと。
- 針刺し損傷が起きた場合は、施設のプロトコルに従い、直ちに適切な処置を取ること。
- 採血中に採血針がホルダーから外れた場合、直ちに患者から抜針し、血液の接触を避け、感染防止措置を取ること。
- ルアーアダプターのルアー接続部は国際規格のルアーフィッティング、ロックフィッティングで規定されている規格に準拠しているが、接続相手が同様の規格に準拠している場合でも、締め方や接続後の取り扱い等により、接続が緩む場合が想定される。確実にしっかりと接続し、漏れ等の異常がないか必ず確認すること。

### <貯蔵・保管方法及び使用期間等>

#### <貯蔵・保管方法>

水ぬれに注意し、直射日光及び高温多湿を避けて保管すること。

#### <使用期限>

箱に記載。

### \* <包装>

- 1箱 48本入り
- 1箱 50本入り
- 1箱 100本入り
- 1箱 200本入り
- 1箱 1000本入り

### \* <製造販売業者及び製造業者の氏名又は名称及び住所等>

製造販売業者：

日本ベクトン・ディッキンソン株式会社  
〒960-2152 福島県福島市土船字五反田1番地  
TEL：0120-8555-90 (BDお客様情報センター)

外国製造業者：

ベクトン・ディッキンソンアンドカンパニー  
(Becton, Dickinson and Company)  
国名：アメリカ合衆国、イギリス、シンガポール、メキシコ



## 機械器具 56 採血又は輸血用器具

管理医療機器 単回使用採血用針 35209002

# BDバキュテイナシステム

## (BD バキュテイナ シングルユースホルダー プラス)


## 再使用禁止

## 【禁忌・禁止】

- 再使用禁止
- 耐圧性能を有するゴムスリーブ付き採血針との組み合わせ以外では使用しないこと。
- 採血終了後、採血管に採血針が刺さったままの状態で駆血帯を外さないこと。[駆血帯を外すことによる圧力の変動により、採血管内の内容物等が患者の体内に逆流するおそれがある。]
- ホルダーは患者ごとの使用とし、使用後は廃棄すること。[ホルダーに血液が付着した場合は、交差感染のおそれがある。]

## 【形状、構造および原理等】

本品は、採血針等を装着し、採血時に採血管を保持する単回使用の器具である。



## 【使用目的、効能又は効果】

真空採血システムの一構成をなすものであり、採血針等を穿刺する際に、固定及び安定化させる用具である。採血針を吻合し、固定した後、当該ホルダー内に真空採血管を挿入することによって、採血が開始される。

## 【品目仕様等】

JIS T 3220 滅菌済み採血用針 6.5.8 ホルダに適合している。

## 【操作方法又は使用方法等】

- 1) あらかじめ手袋を着用する。
- 2) 採血針等をホルダーにセットする。
- 3) 駆血帯を装着し、穿刺部位を消毒する。
- 4) 採血針を血管に穿刺する。
- 5) 採血管を上図に示された位置までホルダーにまっすぐ押し込む。
- 6) 血液の採血管への流入が停止したら、直ちに採血管をホルダーからまっすぐ外す。
- 7) 連続採血する場合には、ホルダーを保持したまま、採血管を取り替える。
- 8) 採血終了後、採血管をホルダーから外した後に駆血帯を外す。
- 9) 抜針し、採血針等をホルダーから外さず適切な耐貫通性廃棄容器に廃棄する。

## ＜使用方法に関連する使用上の注意＞

- 採血針等をホルダーに確実に固定すること。[ホルダーへの固

定が確実でない場合、採血針等が使用中に外れる場合がある。]

- 採血管をホルダーに挿入する際は、ゆっくり押し込むこと。
- フィルム栓タイプ採血管は、採血管先端を図のAのラインに合わせる。オーバーキャップタイプ採血管、ゴム栓タイプ採血管の場合は、キャップが自然に止まる位置に固定して採血すること。それ以上は進めず、採血管先端はBのラインを決して超えないようにすること。
- 採血管を無理に奥まで押し込まないこと。[採血管がホルダーから抜けなくなったり、フィルム栓が破損するおそれがある。]

## 【使用上の注意】

## ＜重要な基本的注意＞

- 併用する医療機器の添付文書を確認後、使用すること。
- 包装が損傷・汚染している場合や製品に破損等の異常が認められる場合は使用しないこと。
- 採血針等のホルダーへの装着時及び装着後に採血針等とホルダーの嵌合が緩い場合には、採血針等及び本品を交換すること。[採血時に採血針がぐらつくおそれがある。]
- 採血針等をホルダーに装着した後はホルダー内に指を入れないこと。[針刺し損傷及び感染のおそれがある。]
- 採血時及び採血管を取り扱う際は、血液との接触を最小限にするため、手袋の着用等の適切な標準予防策をとること。

## ＜併用医療機器＞

- キャップの直径が17mm以上のオーバーキャップタイプの採血管は本品には適合しないため、使用しないこと。[無理に使用すると採血管がホルダーから抜けなくなるおそれがある。]
- BDバキュテイナ採血管のヘモガードタイプキャップ付きの16 x 100mmサイズの採血管は本品には適合しないため、使用しないこと。[無理に使用すると採血管がホルダーから抜けなくなるおそれがある。]

## 【貯蔵・保管方法及び使用期間等】

## ＜貯蔵・保管方法＞

水ぬれに注意し、直射日光および高温多湿を避けて保管すること。

## 【包装】

1袋100個入り

## 【製造販売業者および製造業者の氏名又は名称及び住所等】

製造販売業者：

日本ベクトン・ディッキンソン株式会社  
〒960-2152 福島県福島市土船字五反田1番地  
TEL：0120-8555-90 (BDお客様情報センター)

外国製造業者：

Meiban Micro Pte Ltd  
(メイバン マイクロ ピーティーイー リミテッド)  
国名：シンガポール





---

**Clinical Study Protocol Amendment No 2 Appendix A**

Drug Substance	Dapagliflozin
Study Code	D1692C00012
Edition Number	1
Date	
Protocol Dated	

---

---

**Appendix A**  
**Signatures**

---

## **ASTRAZENECA SIGNATURE(S)**

---

**A long term open label study to evaluate the safety and efficacy of dapagliflozin as monotherapy or combination therapies with anti-diabetic drugs in Japanese subjects with type 2 diabetes who have inadequate glycemic control**

---

This Clinical Study Protocol and all Amendments to the CSP have been subjected to an internal AstraZeneca peer review.

I agree to the terms of this study protocol/amendment.

**AstraZeneca Research and Development  
site representative**

This document contains confidential information, which should not be copied, referred to, released or published without written approval from AstraZeneca. Investigators are cautioned that the information in this protocol may be subject to change and revision.

## **ASTRAZENECA SIGNATURE(S)**

---

**A long term open label study to evaluate the safety and efficacy of dapagliflozin as monotherapy or combination therapies with anti-diabetic drugs in Japanese subjects with type 2 diabetes who have inadequate glycemic control**

---

This Clinical Study Protocol and all Amendments to the CSP have been subjected to an internal AstraZeneca peer review.

I agree to the terms of this study protocol/amendment.

**AstraZeneca Research and Development  
site representative**

This document contains confidential information, which should not be copied, referred to, released or published without written approval from AstraZeneca. Investigators are cautioned that the information in this protocol may be subject to change and revision.



## **ASTRAZENECA SIGNATURE(S)**

---

**A long term open label study to evaluate the safety and efficacy of dapagliflozin as monotherapy or combination therapies with anti-diabetic drugs in Japanese subjects with type 2 diabetes who have inadequate glycemic control**

---

This Clinical Study Protocol and all Amendments to the CSP have been subjected to an internal AstraZeneca peer review.

I agree to the terms of this study protocol/amendment.

**AstraZeneca Research and  
Development site representative**

This document contains confidential information, which should not be copied, referred to, released or published without written approval from AstraZeneca. Investigators are cautioned that the information in this protocol may be subject to change and revision.

Clinical Study Protocol Amendment No 2 Appendix A  
Drug Substance Dapagliflozin  
Study Code D1692C00012  
Edition Number 1  
Date

## **ASTRAZENECA SIGNATURE(S)**

---

**A long term open label study to evaluate the safety and efficacy of dapagliflozin as monotherapy or combination therapies with anti-diabetic drugs in Japanese subjects with type 2 diabetes who have inadequate glycemic control**

---

This Clinical Study Protocol and all Amendments to the CSP have been subjected to an internal AstraZeneca peer review.

I agree to the terms of this study protocol/amendment.

**AstraZeneca Research and  
Development site representative**

This document contains confidential information, which should not be copied, referred to, released or published without written approval from AstraZeneca. Investigators are cautioned that the information in this protocol may be subject to change and revision.